

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/00	A2	(11) International Publication Number: WO 99/61656 (43) International Publication Date: 2 December 1999 (02.12.99)
(21) International Application Number: PCT/US99/11918 (22) International Filing Date: 28 May 1999 (28.05.99) (30) Priority Data: 60/087,170 29 May 1998 (29.05.98) US 60/129,023 13 April 1999 (13.04.99) US (71) Applicant: WISCONSIN ALUMNI RESEARCH FOUNDATION [US/US]; 614 North Walnut Street, P.O. Box 7365, Madison, WI 53707-7365 (US). (72) Inventors: KIMBLE, Judith, E.; 2804 Columbia Road, Madison, WI 53705 (US). BLELLOCH, Robert, H.; 2130 Rusk Street, Madison, WI 53704 (US). (74) Agent: BERSON, Bennett, J.; Quarles & Brady LLP, P.O. Box 2113, Madison, WI 53701-2113 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: AGENT AND METHOD FOR MODULATION OF CELL MIGRATION		
(57) Abstract A GON-1 migration protein in <i>C. elegans</i> and a <i>gon-1</i> gene encoding same are disclosed. The protein, termed GON-1, shows structural similarity to a protein produced by an up-regulated RNA in an advanced tumor cell. Although the tumor cell protein has not previously been identified as having any role in cell migration, it is disclosed herein that the related GON-1 protein is required for cell migration and is involved in shaping tissues or organs. It is deduced that the protein is also a target for modulators of cell migration and tissue shaping.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

AGENT AND METHOD FOR MODULATION OF CELL MIGRATION

5

10

BACKGROUND OF THE INVENTION

Cell migration, particularly migration of cancerous cells and nerve cells, is not well understood, nor are the factors that affect cell migration and tissue shaping in vivo. There is a need in the art to identify and exploit
15 such factors, including but not limited to those involved in normal or abnormal organogenesis. The art also lacks efficient systems for evaluating therapeutic modulators of such functions in vivo and lacks diagnostic methods for assessing the ability of a cell or cell mass to migrate in
20 vivo.

Organogenesis processes in vertebrates proceed in a manner similar to those observed in the common laboratory nematode *C. elegans*. As such, the generation of *C. elegans* gonadal structures can serve as a simple system for
25 investigating developmental morphogenetic processes shared by higher and lower organisms.

In one common morphogenetic process, a tissue bud extends to form an elongate tube with a proximal to distal axis. An emerging theme in bud extension is the presence
30 of specialized regulatory cells at the bud tip that govern elongation. In vertebrate development, this process is

seen in extension of the limb (Johnson and Tabin, 1997; Martin, 1998), ureter (Vainio and Muller, 1997), and lung branches (Hogan, 1998). In the *C. elegans* gonad, long "arms" develop by elongation of buds originating from a gonadal primordium. Each gonadal arm possesses a single "leader cell" that serves this regulatory role (Kimble and White, 1981). The biology of distal tip cell migration during gonadogenesis is known to one skilled in the art of *C. elegans* developmental biology. Indeed, the *C. elegans* gonadal leader cells are among the best defined cells that regulate bud elongation, and therefore serve as a paradigm for investigating this common morphogenetic process.

A second common morphogenetic process of organogenesis is the formation of a complex, differentiated epithelial tube. Formation of a complex epithelial tube can involve an initial condensation of mesenchymal cells, followed by epithelialization, lumen formation, and differentiation into modular units. Vertebrate examples include the kidney tubules (Vainio and Muller, 1997) and heart tube (Fishman and Olson, 1997). Similarly, during *C. elegans* gonadogenesis, cells coalesce to form a compact larval structure called the somatic gonadal primordium (SGP). Following formation of this primordium, cell division and differentiation are accompanied by epithelialization and lumen formation to form a complex tube composed of distinct modular units: the uterus, spermathecae and sheaths in hermaphrodites, and the seminal vesicle and vas deferens in males (Kimble and Hirsh, 1979).

Previous studies have identified several genes in *C. elegans* that influence gonadal morphogenesis. One group of such genes includes *unc-5*, *unc-6*, and *unc-40*, which control the direction of leader cell migration (Hedgecock et al, 1990). Normally, leader cells migrate in one direction, then move dorsally, and finally move in the opposite direction to generate a reflexed gonadal arm. In the absence of *unc-5*, *unc-6*, or *unc-40*, the leader cells fail to turn dorsally. Another gene, *ced-5*, causes the

leader cell to makes extra turns or stop prematurely (Wu and Horvitz, 1998). Therefore, in these mutants, the leader cells migrate, but do not navigate correctly, which results in a failure of the gonadal arms to acquire their normal U-shape. In addition to these genes, others are required for specification of cell fates and also influence morphogenesis (*lin-12*: Greenwald et al., 1983, Newman et al., 1995; *lin-17*: Sternberg and Horvitz, 1988; *lag-2*: Lambie and Kimble, 1991; *ceh-18*: Greenstein et al., 1994, Rose et al., 1997; *lin-26*: den Boer et al., 1998).

A known *C. elegans* genetic locus, *gon-1*, defined by one or more mutants, is essential for extension of gonadal germline arms, but is not responsible for signaling the germline to proliferate. In *C. elegans* hermaphrodites, *GON-1* is required for migration of two distal tip cells to produce two elongated tubes, whereas in males, *gon-1* activity is required for migration of a single linker cell to produce a single elongated tube. In *gon-1* mutant hermaphrodites, the leader cells are born normally in the somatic gonadal cell lineage and function normally to promote germline proliferation, but they fail to migrate and do not support arm extension. Similarly in males, the leader cell does not move and no arm extension occurs. The *gon-1* locus has not heretofore been mapped with particularity to a nucleic acid coding sequence.

Clarification of the genetic basis for *C. elegans gon-1* activity would permit one to apply molecular tools to the study of cell migration in a convenient system. It would be particularly advantageous to find that the *gon-1* locus encodes a protein having structural relationship to proteins of species that are not readily studied in the laboratory, since one would be able to evaluate those proteins in the convenient *C. elegans* system. Such a system would also provide a means for evaluating agents that can modulate the activity of such genes and proteins and would both facilitate understanding the factors involved in cell migration.

BRIEF SUMMARY OF THE INVENTION

In one aspect, the invention can be an isolated polynucleotide coding sequence that encodes a protein that includes both a metalloprotease domain and at least one thrombospondin type 1 domain, where the protein can direct either cell migration or tissue shaping in an analytical system in a target organism as disclosed herein. In another aspect, the invention can also be a variant of the isolated polynucleotide coding sequence that encodes a protein that shares at least 20%, more preferably 50%, still more preferably 70% and most preferably 80% amino acid sequence identity (using GCG Pileup program) with any of the foregoing in the metalloprotease and thrombospondin type 1 domains while also comprising the amino acids of those domains known to those skilled in the art to be required for protein activity. A suitable variant polynucleotide can hybridize under stringent hybridization conditions known to those skilled in the art to a polynucleotide sequence that encodes a protein that can direct cell migration or tissue shaping in the target organism. In one embodiment, a variant polynucleotide can hybridize under stringent hybridization conditions to a *C. elegans gon-1* coding sequence. The variant polynucleotide sequence can be a polynucleotide obtained from an organism or can be a mutated version of any polynucleotide sequence noted above. The variant polynucleotide can encode a protein that is identical or altered relative to the wild-type *C. elegans* GON-1 protein. The encoded protein can have enhanced or reduced activity *in vivo* relative to GON-1.

30 In a related aspect, a polynucleotide coding sequence that encodes a protein having structural and functional similarity with a wild-type or altered migration or shaping protein can also be substituted, in whole or in part, with structurally related or unrelated sequences to encode a heterologous protein or a chimeric protein in the disclosed system, as detailed below.

Applicants herein disclose that the *Caenorhabditis elegans gon-1* activity is encoded by a polynucleotide coding sequence (*gon-1*; SEQ ID NO:1) that encodes an essential protein (GON-1; SEQ ID NO:2) that directs
5 migration of a growing gonadal tube through surrounding basement membranes during gonadogenesis in the nematode and also controls gonadal shape and organ localization.

The migration directing ability and tissue shaping ability are separable and depend upon whether the *gon-1*
10 coding sequence is expressed in distal tip cells or in muscle cells, respectively. In wild-type *C. elegans*, a gonad of normal shape is produced when *gon-1* is expressed in both cell types. Accordingly, one aspect of the invention can also a method for shaping a tissue by
15 selectively expressing a protein associated with both tissue elongation and tissue expansion. GON-1 shares significant amino acid identity with proteins that have been noted in other species.

In a related aspect, the invention can be an isolated
20 and substantially purified preparation of a GON-1 protein, an altered GON-1 protein, a heterologous protein, a chimeric protein, or a variant thereof (referred to herein as "an MPT protein", for reasons discussed below), which can be a target for *in vivo* screening of putative
25 therapeutic modulators, or can be assayed in a diagnostic method for assessing the ability of a cell or cell mass to migrate *in vivo*, or can be exploited as a therapeutic agent to modulate (increase or decrease) *in vivo* cell migration.

One skilled in the art will appreciate that the
30 nucleotide coding sequences and encoded amino acid sequences that fall within the scope of the invention are also subject to natural variation or intentional manipulation (e.g., changes in the nucleotide or amino acid sequence) in ways that do not affect the ability to
35 function as described herein. One skilled in that art also understands that the applicants cannot provide a complete list of nucleotide coding sequences and amino acid

sequences that can function in the methods of the invention. However, in view of the high level of understanding in the art about the amino acids required for activity of proteins that comprise a metalloprotease domain
5 and proteins that comprise a thrombospondin domain, applicants maintain that a skilled artisan can readily determine whether a protein contains both domains. Stöcker, W. et al., "The metzincins - Topological and sequential relations between the atacins, adamalysins,
10 serralyins, and matrixins (collagenases) define a superfamily of zinc-peptidases," Protein Science 4:823-840 (1995), Rawlings, N.D. and A.J. Barrett, "Evolutionary families of metallopeptidases, Methods in Enzymology 248:183-228 (1995), and Adams, J.C. et al., The
15 Thrombospondin Gene Family, R.G. Landes Company, Austin, TX (1995), all incorporated herein by reference in their entirety, provide sufficient guidance to permit those in the art to establish whether a protein comprises both a metalloprotease and a thrombospondin domain.

20 The invention is further summarized in that an antibody can be produced against characteristic epitopes of any of the foregoing proteins using standard methods. The antibody can be used both diagnostically to ascertain the presence of an MPT protein, or therapeutically to interfere
25 with activity of the MPT protein.

The present invention is also summarized in that an animal that contains a *gon-1* allele (or homolog or variant thereof) is a convenient screening tool for finding modulators of cell migration. The present invention is
30 thus further summarized in that a method for identifying modulators of the disclosed MPT proteins includes the steps of treating a target organism having a cell that can migrate or be shaped when under control of an MPT protein with at least one potential modulator of migration or
35 shaping and observing in the treated target organism a change in migration or shaping of the cell or tissue attributable to the presence of a modulator. In a

preferred embodiment, the cell is a developing gonadal cell in *C. elegans*, although other cells or organs may be similarly regulated by MPT proteins in other organisms.

The ability of the MPT protein to direct a cell or
5 tissue under its influence to migrate or be shaped can be modulated (increased or decreased) in a variety of ways, such as by altering the migration protein's primary, secondary, or tertiary structure, by altering the location or amount of the protein in an organism, by altering the
10 transcriptional or translational regulation of the gene that encodes the protein, or by providing the organism with an agonist or antagonist molecule in an amount sufficient to interact with the MPT protein so as to increase or decrease the ability of the protein to direct migration or
15 shaping.

In a related method, one can also identify nucleic acid sequences required or desired for migration or shaping of such a cell, by treating a target organism with an agent that affects the polynucleotide sequences of the target
20 organism that encode the MPT protein or that participate in regulating expression of the MPT protein, and then identifying sequences affected by the treatment. The sequences identified in the method can be either complete or partial coding sequences or can be regulatory sequences.

25 It is an object of the present invention to identify a protein and nucleotide sequence encoding same that directs migration or shaping of a cell or tissue.

It is another object of the present invention to provide a method for modulating cell migration or shaping.

30 It is yet another object of the present invention to provide a system and method for screening putative modulators of migration or shaping of cells or tissues.

It is an advantage of the present invention that agents having a putative effect upon migration or shaping
35 can be screened in a convenient model system rather than in a vertebrate organism.

Other objects, features and advantages of present

invention will become apparent upon consideration of the following detailed description taken in conjunction with the accompanying drawings.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

5 Fig. 1A depicts a schematic map of the *gon-1* locus in *C. elegans* from which the gene was cloned and shows the exon-intron structure of *gon-1*.

Fig. 1B shows a schematic map of *C. elegans* GON-1, the location of five protein-truncating stop mutants in GON-1
10 and a comparison to the protein structures of the murine ADAMTS-1 protein, and the bovine procollagen-I N-proteinase (PN1P) protein. From left to right, GON-1 includes a prodomain, a metalloprotease domain, a first cysteine rich region, a thrombospondin type I motif, a second cysteine
15 rich region, and a plurality of thrombospondin type I-like motifs. The five mutants are identified as *q518* (aa591 TGG->TGA), *e2551* (aa1069 TGG->TAG), *e2547* (aa1229 TGG->TGA), *q18* (aa1234 TGG->TAG) W->stop, and *e1254* (aa1345 CGA->TGA) R->stop).

20 Fig. 1C compares the *C. elegans* GON-1 amino acid sequence to sequences of the ADAMTS-1 and PN1P proteins. In the metalloprotease domain, amino acids important for enzymatic activity are marked by an asterisk (*). Three conserved histidines (GON-1, aa 424, 428, 434) bind a
25 catalytically essential Zn^{+2} ion in well characterized metalloproteases, while a glutamic acid residue (GON-1, aa 425) is thought to be directly involved in cleavage (Stöcker et al, 1995). In addition, two conserved glycines and a downstream methionine seem to be important for
30 structure of the active site. GON-1 bears one of the glycines (aa 427) and the methionine (aa 454), but the second glycine is changed to serine in GON-1 (aa431). In the canonical TSPT1 domain, amino acids conserved in vertebrate TSP type-1 repeats are shown by a plus (+). The
35 mutation, *gon-1(q518)*, is marked by an inverted triangle

(V). For the TSPT1-like repeats, only 2 of the 17 are shown. The consensus sequence for these repeats is:
W-X₄₋₅-W-X₂-CS-X₂-CG-X₄₋₅-X-G-X₃-R-X₃-C-X₄₋₂₇-C-X₈₋₁₂-C-X₃₋₄-C.
Because only the first two TSPT1-like motifs are shown, the
5 other mutations are not indicated in this figure.

Fig. 2A depicts normal morphogenesis of the *C. elegans* hermaphrodite gonad.

Fig. 2B shows that arm extension does not occur in *gon-1* mutants and that the gonad develops as a disorganized
10 mass of somatic and germline tissues. Similarly, in males, the *gon-1* mutant gonad is severely disorganized and does not acquire its normal shape.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The existence of a protein in *C. elegans* required for
15 cell migration or shaping has not heretofore been known, nor has any function been previously ascribed to a protein encoded by the designated sequence. The inventors have determined that a functional GON-1 protein is required for migration of the regulatory cells that lead the developing
20 gonad organ during its migration. GON-1 is also involved in shaping tissues such as gonads. By appreciating the role of GON-1 (and the *gon-1* gene) and its relationship to a related gene that is upregulated in a metastatic tumor cell, the inventors have identified a gene and protein
25 believed to be fundamental in the process of normal and abnormal cell migration and tissue shaping. The gene and protein, and related genes and proteins, can be utilized in the methods of the invention as described herein.

References herein to influencing cell migration are also
30 intended to encompass shaping of tissues or organs. Likewise, references to a migration protein encompass proteins of the same class that can also be used in methods for shaping tissues or organs.

Generally speaking, the methods of the present
35 invention permit one to identify agents that modulate cell migration or tissue shaping *in vivo* or *in vitro*. One can

treat target organisms with panels of polynucleotides, proteins, sugars, lipids, organic molecules, other chemicals, synthetic or natural pharmaceutical agents or other agents to determine whether any agent affects
5 activity of an MST protein. This list is necessarily incomplete, since one cannot predict in advance which agents will be effective. However, applicants have enabled a system for screening panels of putative agents, in accord with the common practices of pharmaceutical companies that
10 typically screen thousands of compounds against a test system in an effort to reveal preferred agents. Candidate agents likely to modulate MPT proteins in the disclosed system include tissue inhibitors of metalloproteases and pharmaceutical metalloprotease inhibitors or enhancers such
15 as those from British Biotech. Inhibitors or enhancers of thrombospondin activity are also good candidate agents.

Agents so identified can be used therapeutically to enhance or inhibit cell migration or to influence tissue shape. Agents having an adverse or inhibiting or knock-out
20 effect upon activity of a migration protein can also be used in a method for biocontrol of animals that employ the migration protein in gonadal development, where the method includes the step of exposing a developing animal to an amount of the agent effective to prevent gonadal
25 development such that the animals are rendered sterile. While this biocontrol method is particularly envisioned for use in nematodes, it may be applicable to other animals as well, since genes related structurally and functionally to *gon-1* are known to exist in animals as diverse as
30 nematodes, cattle and humans.

Using the invention one can also identify polynucleotide sequences including coding and regulatory sequences that affect activity of a migration protein. For example, null or so-called reduced activity mutants can be
35 mutagenized and assayed for activity-restoring, activity-inhibiting or activity-enhancing changes. By extension, one can perform comparable screens *ad infinitum* on

sequences identified in this manner, to obtain still more sequences that have an indirect effect on migration activity. After identifying such sequences in a target organism, one can obtain homologous polynucleotides from
5 other organisms by screening nucleic acid libraries under stringent hybridization conditions in a manner known to those skilled in the art.

A method for evaluating putative modulators of cell migration preferably employs a nematode as a target
10 organism. The methods may be advantageously practiced using a nematode that comprises a migration protein as described herein, or a mutant nematode that either lacks a migration protein or contains a migration protein having reduced activity. The protein can be encoded by wild-type
15 *C. elegans gon-1* (disclosed herein), by a mutant that confers upon the nematode an enhanced or reduced sensitivity to modulators, by a transgene from another organism, in whole or in part, or by a variant of any of the foregoing. Nematodes are desirable target organisms,
20 in general, because they are easy to grow and maintain, and easy to assay, particularly because they are transparent.

Nematodes are also particularly desired because the powerful techniques of reverse genetics can be employed. One can also target specific *C. elegans* sequences for
25 mutation or RNA-mediated interference (a technique used to transiently knock genes out by RNA injection) to identify nucleic acid and protein sequences that have a direct inhibitory or enhancing effect on *gon-1* activity.

With the identification of the *gon-1* gene and GON-1
30 protein in *C. elegans* and the discovery of homologous genes in other species, the functions of migration proteins can be analyzed *in vivo* during organogenesis using the full force of molecular genetics available in that system. Such functions can include, but may not be limited to cell
35 migration, basement membrane remodeling, and tubular organ formation.

Although the system is exemplified in *C. elegans*, a

free-living (i.e., non-parasitic) nematode, those skilled in the art can develop similar systems operating on the same principles without undue experimentation in other convenient organisms, including other nematodes including, 5 without limitation, *C. briggsae*, or in, for example, *Drosophila*, or other organisms conveniently studied in the laboratory. To do so, one would only need to identify the homolog of *gon-1* in such an organism, using standard molecular biological methods and then screen for related 10 genes, proteins and other factors as described herein. One could also use such systems in other animals to study transgenes in ways comparable to those described herein. Those skilled in the art can produce transgenic animals of many species without undue experimentation.

15 In the method, a putative modulator is provided to the target organism, for example, by adding it to the growth media, by injecting it into the organism or by gene transformation technology. The effects of said modulator can be assessed either by screening for changes in cell 20 migration or by genetic selection for fertile animals. The assessment methods are known to those skilled in the art. *Caenorhabditis elegans*: Modern Biological Analysis of an Organism, Methods in Cell Biology, volume 48, Epstein, H. F. and D. C. Shakes, eds., Academic Press (1995), 25 incorporated herein by reference in its entirety, describes suitable methods and conditions for growing and monitoring *C. elegans*.

C. elegans GON-1 is characterized by a multi-domain structure that includes several known motifs. GON-1 protein 30 is a secreted metalloproteinase that lacks a transmembrane domain and possesses a predicted metalloprotease domain between amino acids 269-456. The metalloprotease enzymatic activity is essential for GON-1 function; proteins that might be cleaved by this metalloproteinase include 35 components of the basement membrane and other proteins that modulate migration. The metalloprotease domain shares sequence similarity with other metalloproteinase enzymes.

In addition to its metalloprotease domain, GON-1 possesses a series of consecutive motifs that are related to, but variants of, the thrombospondin type 1 (TSPT1) repeats (Fig. 1B,C). The most N-terminal TSPT1 repeat bears the hallmarks of this type of motif in vertebrate thrombospondins (15/16 of the consensus amino acids, + in Fig. 1C) (Adams et al., 1995), whereas the remaining 17 repeats are less similar and define a TSPT1-like variant. Proteins that might interact with this domain include proteins that modulate migration, including but not limited to components of the basement membrane.

GON-1 is similar to members of the reprotolysin subfamily (Rawlings, N.D. and A.J. Barrett, "Evolutionary families of metalloproteases, Methods in Enzymology 248:183-228 (1995), incorporated herein by reference in its entirety). At the N-terminal border of the metalloprotease domain, there is a potential furin cleavage site (Fig. 1C) (Pei and Weiss, 1995; Pei and Weiss, 1996). GON-1 and the reprotolysins share a common zinc binding active site with the larger metzincin superfamily (Stöcker et al., 1995). Amino acid conservation within the active site together with the known crystal structure of several superfamily members reveals those amino acids essential for enzymatic activity (marked by asterisks in Fig. 1c) (ibid). GON-1 has all amino acids implicated in catalysis and all but one implicated in structure of the active site.

Wild-type *C. elegans* GON-1 (SEQ ID NO:2) is suitable for use in the methods of the present invention, although a skilled artisan can replace the *C. elegans gon-1* coding sequence with a sequence that encodes all or part of a homologous protein, using the standard tools available to a molecular biologist. This mixing and matching can increase or decrease the activity of the encoded chimeric protein. As described elsewhere herein, it can be desirable to provide a system having reduced or enhanced migration activity, or even no migration activity, depending upon whether one is evaluating agents that enhance or inhibit

migration. Increased gene activity is characterized either by increased gonadal arm extension, increased compactness of gonadal tissue, or fertility. Decreased gene activity is assayed either by decreased gonadal arm extension,
5 decreased compactness of gonadal tissue or sterility. Certain specific activity-reducing mutations in *gon-1* are described in the Examples.

Sequences with related structures have already been isolated from vertebrate organisms, but no related
10 invertebrate sequence is known to the inventors. Still other related metalloprotease proteins (and polynucleotide sequences encoding same) will be isolated from vertebrate and invertebrate organisms. While the *C. elegans gon-1* protein includes 17 thrombospondin domains, the bovine and
15 murine homologs include only 2 such domains. Other known members of the family also have one canonical TSPT1 repeat, can contain at least one TSPT1-like variant repeat, and contain two conserved cysteine rich regions. Based on this conserved architecture, we suggest the name MPT (for
20 MetalloProtease with TSP1 repeats) for the family.

While the *in vivo* functions of these proteins may differ from that of *C. elegans GON-1*, these proteins are expected to function in place of *GON-1* in whole or in part in the disclosed methods. All such homologs from other
25 vertebrate and invertebrate organisms (and the polynucleotide sequences that encode such homologs), variants thereof, and chimerics that incorporate portions thereof, whether obtained naturally or induced in the laboratory using the tools available to a molecular
30 biologist, are considered to be useful in the present invention. In particular, functional domains, such as the metalloprotease domain, can be swapped into corresponding domains in *gon-1*.

The amino acid sequences of *GON-1*, *ADAMTS-1* and bovine
35 *PN1P* are compared in Fig. 1C. The additional thrombospondin domains of *GON-1* not found in *ADAMTS-1* or *PN1P* are not shown in Fig. 1C. Those portions of *GON-1*

that have no obvious relationship to known motifs are conserved among the family of GON-1 homologs. The GON-1 protein shows significant sequence similarity to the bovine procollagen-1 N-proteinase (P1NP), to the murine ADAMTS-1
5 protein, and to a pair of human aggrecan-degrading metalloprotease-encoding sequences described in International Patent Application Number PCT/US98/15438, published on February 4, 1999 as International Publication No. WO 99/05291, incorporated herein by reference in its
10 entirety. Another human homolog which has significant identity to the bovine P1NP has Genbank accession number d1021662.

Bovine P1NP can proteolyze the N-terminal propeptide from collagen I (Colige et al., 1995, Colige et al., 1997).
15 Metalloprotease activity is required for GON-1 function and suggest that, like P1NP, it may cleave components of the extracellular matrix. Murine adamts-1 expression correlates with tumor cell progression (Kuno et al., 1997). The murine ADAMTS-1 protein is found in an advanced
20 cachexogenic murine tumor cell. Human aggrecanase has been associated with arthritis in humans. Given the role of GON-1 in regulating cell migration of the *C. elegans* leader cell, we suggest that MPT proteins may be involved more generally in cell migrations that must pass through
25 extracellular matrix and that, in cancerous tissues, loss of MPT regulation may promote metastasis. The percent identity of the identified domains of *C. elegans* GON-1 with the bovine and murine proteins is shown in Fig. 1B.

Changes can be made in any of the foregoing at the
30 nucleic acid level in a manner known to those skilled in the art, by, for example, removing a section of the coding sequence, interrupting the coding sequence with an additional sequence, rearranging at least one section of the gene, or by providing in the sequence other changes
35 that can include but are not limited to point mutations that either truncate the protein or disable an active site in the protein encoded by the altered polynucleotide.

Changes can also be made by altering the transcription or translation of the gene that encodes the migration protein by altering in a manner known to the art the upstream and/or downstream regulatory sequences that the
5 surround the gene. Likewise the translation-regulating elements of an mRNA encoding the migration protein can also be altered to affect the stability or location of the mRNA. An antisense RNA can also interfere with translation of the migration protein.

10 At the protein level, one skilled in the art can modulate the activity of the migration protein either by modifying the protein encoded by the gene as noted above or by directing the protein to be modified in vivo, for example, by providing in the protein appropriate signal or
15 signals for cleavage or degradation by other cellular factors. Alternatively, the protein can be targeted with an activity-modulating factor such as a protein, a peptide, or an organic or inorganic co-factor. Any of these factors can, for example, occupy or obstruct an active site of the
20 protein which is required for activity. Likewise, if the activity of the protein is natively regulated by an endogenous co-factor, an effect can be achieved by modulating the availability of the native co-factor.

One skilled in art is familiar with the techniques
25 associated with the aforementioned alterations, including the production of any construct necessary to effect such changes. One skilled in the art also understands that changes in the primary amino acid sequence (including, e.g., substitutions, deletions, additions, inversions) may
30 or may not alter the activity of a protein, depending upon the position and the extent of the change.

For purposes of this application a migration protein is considered active if it causes a cell that comprises the protein, or a cell that is under the influence of the
35 protein, to migrate to any appreciable extent. A cell is "under the influence of the protein" if the cell migrates in the presence of the protein, even if the cell does not

contain the protein. *In vivo*, the cell from which the protein is secreted and its site of action remain unknown.

Non-native transgene sequences containing non-native sequences homologous to all or part of *C. elegans gon-1* can be introduced into *C. elegans* on an expressible genetic construct that contains a promoter that drives expression in a tissue that allows easy assay so that the effect or effects of those sequences on migration and other functions can be evaluated in the system. Methods for generating and selecting transgenic nematodes are well-known in the art. Transgenic animals can rescue null mutants or can suppress or enhance the activity in the reduced-activity mutants. A preferred example of a transgene sequence is a human *gon-1* homolog sequence, although any of homolog can be used. Some constructs may contain all or part of the *gon-1* coding sequences. The transgene should be appropriately expressed near the cells to be controlled by the migration protein. In *C. elegans*, the *gon-1* promoter, active in leader cells and in muscle cells, is suitable. Other promoters that can be used in *C. elegans* include the *lag-2* promoter, which drives expression in the hermaphrodite distal tip cells, and the *unc-54* promoter which drives expression in body wall muscle.

One can assay for effects of treatment with a potential modulating agent on cell migration and gonadal tube extension by comparing migration after treatment to the cell migration in either a wild-type organism or to that in an untreated, previously characterized mutant. Before treatment in the methods, if the migration protein is expressed in leader cells at wild-type levels, directed elongation of gonadal arms along a proximal-distal axis is observed. If the migration protein is expressed in muscle, on the other hand, one observes more dispersed activity, which may be important for expansion as the gonad along the dorsal-ventral and left-right axes. If a migration protein having a level of activity comparable to that of the wild type protein is expressed from a polynucleotide sequence

under control of the native *gon-1* promoter, of course, normal gonadal development is observed, as is shown in Fig. 2A. Fig. 2B shows that arm extension does not occur in *gon-1* mutants and that the gonad develops as a disorganized mass of somatic and germline tissues. Similarly, in males, the *gon-1* mutant gonad is severely disorganized and does not acquire its normal shape. Both wild-type activity and the mutant phenotype can be modified by treatment according to the methods. One can also direct the shape of a tissue or organ by introducing a transgene coding sequence under control of a promoter selected to express the transgene coding sequence in a desired tissue or cell type.

One can also assess whether a cell has the potential for migration by analyzing for example, the level of the migration protein in the cell, or the level at which the RNA encoding the migration protein is present. A diagnostic assay for the presence of active site residues in the protein can also be devised. Likewise, the presence or absence of a DNA sequence encoding an essential aspect of the protein can also be used in a diagnostic manner to assess the likelihood of cell migration.

Our finding that GON-1 is tightly regulated to achieve arm extension during gonadogenesis in *C. elegans* suggests that similar activities may play similar roles in the morphogenesis of organs throughout the animal kingdom. Previous *in vitro* experiments support this notion. For example, antibodies recognizing matrix metalloprotease 9 (MM9) can block branching of the ureter bud during kidney development (Lelongt et al., 1997), and inhibitors of MMPs block the invasion of endothelium cells into a fibrin matrix in assays for angiogenesis (Hiraoka et al., 1998). Based on these observations and our analysis of GON-1, we suggest that the MPT metalloproteases are critical modulators of organogenesis.

Whether the target organism contains a wild-type *C. elegans gon-1* gene, a mutant *gon-1* gene or a transgene substituted in place of *gon-1*, in whole or in part, the

system is readily used to identify other genes, proteins, drugs, chemicals or other factors that either enhance or antagonize activity.

In a method for increasing the migration of the cell,
5 the native protein or related protein or a genetic
construct encoding same can be administered to, or caused
to be expressed at a high level in, the target cell.
Alternatively, an enhancing factor can be provided inside
or outside the target cell, as appropriate. Where it is
10 desired to decrease migration of a targeted cell, as in the
case of a tumor cell, an inhibiting factor can be added
into, or the vicinity of, the targeted cell. The vicinity
of the cell is defined as sufficiently close to the
targeted cell so as to effect a desired change in the cell
15 migration. If the migration protein is secreted from the
cell in which it is produced, the activity of the protein
can further be modulated either by preventing secretion of
the protein or by interfering with the protein activity
outside the cell. If the protein acts outside the target
20 cell, the protein, an active portion thereof, or a
modulating factor can be administered to the vicinity in an
amount effective to modulate cell migration.

The reproductive sterility that can result from
inhibited migration of developing gonadal cells under the
25 control of an migration protein that is inactive or has
reduced activity can be further exploited, for example, in
a method for controlling reproduction of an organism that
relies upon a migration protein during gonadogenesis. An
organism for which such control would be appropriate would
30 include *C. elegans* and other nematodes or parasites, and
could include other invertebrates, as well as vertebrate
species including, for example, avian, amphibian, reptilian
and mammalian species.

With an appreciation for the migration proteins of the
35 invention, normal and abnormal cell migration attributable
to activity of a migration protein can be therapeutically
increased or decreased. The mechanisms by which the gene

and protein are regulated can be determined by one skilled in the art and can be advantageously exploited to modulate expression of the migration protein at either the nucleic acid or protein levels.

5

EXAMPLES

To gain molecular insight into *gon-1* function, we cloned the gene by a combination of fine genetic mapping, mutant rescue and RNA-mediated interference. Mutations in the *gon-1* gene were finely mapped by genetic crosses with
10 respect to markers that had already been placed on the physical map. Cosmids in the region were next tested for mutant rescue of the *gon-1* mutations. The genomic *C. elegans* sequence that includes the coding sequence of the *gon-1* gene in a plurality of exons is found on cosmids
15 F25H8 (Accession # 69360) and T13H10 (Accession #69361); T13H10 bears most of *gon-1* and rescued the *gon-1* phenotype. The predicted open reading frames on this cosmid were tested by RNA-mediated interference to identify the transcript corresponding to *gon-1* activity. The
20 identification of this transcript as *gon-1* was then confirmed by subcloning and mutant rescue by a smaller region of the cosmid that contained that transcript, by RNA-mediated interference, and by identifying *gon-1* mutations in the coding region of this transcript. The
25 positions in the migration protein that correspond to the identified mutations are indicated in Fig. 1B. We confirmed identification of F25H8.3 as *gon-1* by identifying molecular lesions for a plurality of *gon-1* alleles.

Mutants were obtained as described (Brenner, S. "The
30 Genetics of *Caenorhabditis elegans*, Genetics 77:71-94 (1974), incorporated herein by reference. Each contained an allele of *gon-1* that maps to chromosome IV between *unc-24* and *dpy-20*, all are recessive, and all are fully penetrant for sterility. Five alleles, *e1254*, *e2547*, *q18*,
35 *q517*, and *q518*, fail to complement the sixth allele, *e2551*, and, therefore, the mutations define a single gene. Three-factor mapping places *gon-1*(*e2551*) 0.08 map units to

the right of *elt-1* and 0.12 map units to the left of *unc-43* at position 4.44. Specifically, among *Unc-43* non-*Elt-1* recombinants isolated from *gon-1/ elt-1 unc-43* mothers, 8/13 carried the *gon-1* mutation.

5 To compare allelic strengths, we examined the penetrance of arm extension defects in homozygotes for each allele. In *gon-1(q518)* homozygotes, no arm extension was observed at 15°, 20° or 25°C. However, in homozygotes for the other *gon-1* alleles, some arms extended at least
10 partially. By this measure, the *gon-1* alleles can be placed in an allelic series: *q518* < *e2547* ≈ *q18* < *e1254* ≈ *q517* < *e2551*. Interestingly, the weaker *gon-1* alleles have a more severe defect at lower temperature, which may reflect a cold sensitivity of *GON-1* function, or of the
15 process of arm extension itself.

The strongest loss-of-function allele is *gon-1(q518)* which is a nonsense mutation that resides in the canonical TSP1 motif; the other mutations are located in the TSP1t1-like repeats. *gon-1(q518)*, the nonsense mutant
20 located closest to the N-terminus, has the most severe effect on cell migration; nonsense mutants located closer to the C-terminus than *q518* are partially defective for migration. Because the mutant phenotype for *gon-1(q518)* homozygotes is identical to that of *gon-1(q518)* hemizygotes
25 and because *gon-1(q518)* bears a nonsense mutation predicted to remove the bulk of the *GON-1* protein, this allele is likely to be a molecular null. Therefore, *gon-1(q518)* was used for analyzing the roles of *gon-1* in gonadal morphogenesis and is referred to as *gon-1(0)*.

30 Normally, the gonad is a tubular structure with specialized regions. By contrast, in *gon-1* mutants, the adult gonadal tissues exist as a disorganized mass with little or no tubular morphology. Specifically, neither arms nor somatic gonadal structures (e.g. uterus,
35 spermatheca) are observed. In all cases, however, the gonads are rendered infertile by these mutations.

In *C. elegans*, mRNAs containing premature stop codons

are normally degraded by the *smg* system, but those mRNAs are stabilized in a *smg* mutant background (Anderson and Kimble, 1997). Therefore, the remaining activity of truncated GON-1 proteins should be evident in *smg-1; gon-1* double mutants. We found that *gon-1(q518)* was not suppressed in a *smg* background, whereas all four mutations in the TSP1-like repeats were suppressed. Therefore, while the GON-1(q518) mutant protein that possesses the metalloprotease domain but lacks the *bona fide* TSPT1 motif (as well as the rest of the protein C-terminally), is not capable of mutant rescue, the other truncated proteins are. The conclusion that two TSPT1-like repeats are sufficient for rescuing activity was confirmed by mutant rescue with a mini-transgene.

The lack of gonadal arms in *gon-1(0)* mutants suggested that the leader cells, which normally govern arm extension, may be defective. To assess whether leader cells were generated during development, we first examined the gonadal cell lineages in *gon-1(0)* mutants during the first two larval stages. Normally, the somatic gonadal progenitor cells, Z1 and Z4, give rise to two leader cells, Z1.aa and Z4.pp, in hermaphrodites, and one leader cell, Z1.pa or Z4.aa, in males (Kimble and Hirsh, 1979). In hermaphrodites, these leader cells are called distal tip cells (DTC), and in males, they are called linker cells (LC). The hermaphrodite distal tip cell is both a leader cell and a regulator of germline proliferation. Kimble, J.E. and J.G. White, "On the control of germ cell development in *Caenorhabditis elegans*, Devel. Biol. 81:208-219 (1981), incorporated herein by reference in its entirety, provides guidance for a skilled artisan on the biology of distal tip cell migration. The information disclosed in that paper can be employed in determining whether an agent modulates cell migration or tissue shaping in a method of the invention.

In *gon-1(0)* hermaphrodites and males, we found that the timing and pattern of cell divisions of Z1 and Z4 and

their descendants were the same as in wild-type during L1 and L2 (data not shown). In particular, Z1.aa and Z1.pp in hermaphrodites and Z1.pa/Z4.aa in males were born at the correct time and place. To ask whether the presumptive
5 hermaphrodite leader cells, Z1.aa and Z4.pp, had adopted the leader fate, we examined expression of a molecular marker for that fate. The *unc-5* gene encodes a netrin receptor and is essential for dorsal migration of leader cells (Leung-Hagesteijn et al, 1992). Using a reporter
10 transgene, *unc-5::lacZ* (J. Culotti, personal communication), we found that *unc-5* expression was the same in wild-type and *gon-1(0)* animals: *unc-5* was not expressed during early larval stages, but was activated in late L3 when the DTCs normally turn dorsally during
15 wild-type gonadogenesis.

Since the hermaphrodite leader cells, Z1.aa and Z4.pp, also control germline proliferation, we next asked if they were correctly specified for that regulatory function. To this end, we examined expression of the *lag-2* gene, which
20 encodes the DTC signal for germline proliferation (Henderson et al., 1994). Using a reporter transgene, *lag-2::GFP*, we found that *lag-2::GFP* expression was similar in wild-type and *gon-1* gonads. Furthermore, we ablated Z1.aa and Z4.pp in *gon-1(0)* mutants and found that germline
25 proliferation was arrested. Therefore, the hermaphrodite DTCs, Z1.aa and Z4.pp, appear to be specified correctly both as leader cells and as regulators of germline proliferation.

Since the leader cells appeared to be specified
30 correctly in *gon-1* mutants, we next examined their ability to migrate and lead arm extension. Normally, the hermaphrodite leader cells (distal tip cells) migrate away from the center of the gonad along the anterior-posterior axis, then reflex dorsally, and migrate back. To compare
35 leader cell migration in wild-type and *gon-1(0)* mutants, we followed their movements throughout gonadal development and at the same time measured gonadal lengths. At the

mid-L1 stage, just prior to division of the leader cell progenitors, Z1 and Z4, the length of the gonad from anterior to posterior end was 19 μm in both wild-type and *gon-1(0)* mutants. Following division of Z1 and Z4 in late L1, a small difference in gonadal length was discerned: 25 μm in wild-type vs. 22 μm in *gon-1* mutants. However, in older larvae with differentiated leader cells, the length differences were dramatic. In *gon-1(0)* hermaphrodites, the distal tip cells had moved little from their birth position and little to no gonad extension had occurred.

A similar defect is observed in males. Normally, the male leader cell (linker cell) migrates anteriorly, then reflexes and migrates to posterior end of the worm. However in *gon-1(0)* males, the linker cell failed to migrate, and little to no extension had occurred. We conclude that *gon-1* is required for leader cell migration and hence gonadal arm extension.

As we observed leader cells during gonadogenesis, we noticed that they assumed an unusual morphology. To explore this further, we examined hermaphrodite DTCs using fluorescence and thin section electron microscopy (EM). Using *lag-2::GFP*, which is expressed in hermaphrodite DTCs and reveals the extent of their cytoplasm (D. Gao and J. Kimble, unpublished), we found that the wild-type and *gon-1(0)* DTCs had dramatically different morphologies. In wild-type, the DTC was crescent-shaped with processes extending around the germ line, while in *gon-1* mutants, it was round and enlarged. Furthermore, the position of the nucleus within the DTC was variable in *gon-1* mutants, whereas in wild-type, it was located at the leading edge of the migrating cell. By EM, we confirmed the difference in morphology between wild-type and *gon-1* leader cells and also discovered a difference in subcellular organization. Whereas wild-type leader cells extend processes along the germline, *gon-1(0)* leader cells do not possess such processes. Furthermore, the plasma membrane is abnormally invaginated in *gon-1(0)* L3 leader cells, and these

membranes accumulate within the cytoplasm of older *gon-1(0)* mutants.

The lack of gonadal arms is not the only defect in *gon-1* mutants. In addition, no gonadal structures (e.g. uterus in hermaphrodites, vas deferens in males) can be discerned. One problem might have been a failure to differentiate gonadal tissues. However, we were able to identify the major somatic gonadal cell types in late L4 *gon-1(0)* mutants. To see somatic gonadal sheath cells, we used *lim-7::GFP*, which expresses Green Fluorescent Protein (GFP) in hermaphrodite sheath cells (O. Hobert, pers. comm.). In wild-type, fluorescence from *lim-7::GFP* encircled the germ cells, while in *gon-1* mutants, only irregularly-shaped patches were observed. Similarly, MH27 antibody, which stains spermathecal cells intensely (den Boer et al., 1998), was present in disorganized patches in *gon-1* mutants. Finally, cells with a typically uterine morphology were present, but no normal uterine structure was found in *gon-1* mutants. Therefore, the gonadal tissues in *gon-1(0)* mutants appear to differentiate correctly.

One simple explanation for the gross morphogenetic defects of mature *gon-1* gonads might have been that all aspects of gonadal morphogenesis are disrupted as a consequence of the defect in leader cell migration. Indeed, by killing the distal tip cells in wild-type animals, we could reproduce the *gon-1* mutant phenotype: arms did not extend and gonadal structures were grossly malformed. However, closer inspection suggests that *gon-1* has a role in gonad morphogenesis independent of leader cells.

To examine the generation of gonadal somatic structures, we removed the germ line (-GL) from *gon-1(0)* to permit formation of an essentially normal somatic gonadal primordium at the early L3 stage and we removed both leader cells (-DTCs) and germline (-GL) from wild-type hermaphrodites as a control. The control animals had no arm extension, but formed a normal somatic gonadal primordium.

A comparison of gonadal structures at the L4 stage, when they are most easily scored, revealed striking differences. While fragments of uterus were present in *gon-1(-GL)* hermaphrodites, no coherent uterus was observed.

- 5 Furthermore, the *gon-1 (-GL)* gonad was small, and most gonadal had extruded from the gonad proper. By contrast, an apparently normal uterus formed in the wild-type animals lacking both DTCs and germ line. Therefore, *gon-1* is required not only for arm extension, but also for
10 morphogenesis of the uterus.

Finally, we asked whether *gon-1* functions in the development of non-gonadal tissues. We assayed embryonic viability, the overall shape of the animal, coordination of its movements, mating behavior in males, the male tail,
15 growth rate, and entry and exit into dauer stage of the life cycle: all were normal in *gon-1(0)* mutants. The normal movement and shape of *gon-1(0)* mutants suggests that *gon-1* is not required generally for cell migration. For example, failure in migration of the CAN neuron causes the
20 tail to wither (Forrester et al., 1998), and defects in axon migration leads to an uncoordinated (*Unc*) phenotype (Hedgecock et al., 1990). Furthermore, we followed the M sex myoblast and the Q neuroblasts migrations (Antebi et al., 1997) in at least five *gon-1(0)* mutants, and both were
25 normal. We conclude that *gon-1* does not affect cell migrations generally and, furthermore, that *gon-1* does not affect the development of non-gonadal cells, tissues or organs. Finally, we examined the non-gonadal tissues in *gon-1* mutants that had been operated during L1 to remove
30 Z1-Z4, the four gonadal progenitor cells. This experiment was done, because the disorganized gonadal tissues in *gon-1(0)* hermaphrodites often cause the animal to explode during adulthood, preventing examination of their non-gonadal tissues at this stage. Although these
35 gonadless *gon-1* adults had no gross defects, we observed a reproducible vacuolization in the body wall with differential interference contrast microscopy, which was

not seen in similarly treated wild-type animals. However, it must be emphasized that this defect has no apparent developmental consequences. Given the dramatic effects of *gon-1* on gonadogenesis, we suggest that the major role of *gon-1* in development is to control the shape of the gonad.

The wild-type *C. elegans gon-1* sequence is shown in SEQ. ID. NO. 1. The protein encoded by SEQ. ID. NO. 1 is shown in full in SEQ. ID. NO. 2 and in part in comparative Fig. 1C.

10

PROPHETIC EXAMPLE

A target organism that contains a migration protein is treated with one or more potential modulators of migration of a developing gonadal cell. The organism is preferably a nematode, and is more preferably *C. elegans*. The potential modulating agent is administered in an amount typical of any additive to a culture, preferably at a level of several nanograms to several micrograms per milliliter. The organism can contain a native migration protein or a variant form of a native migration protein, or can express a migration protein from a transgene that can be delivered to the organism in a manner known to those skilled in the art. The protein can also be a chimeric protein expressed from a transgenic polynucleotide that comprises sequences from at least one of the foregoing polynucleotides.

Upon examination, it is observed that one can rescue migration in a target that lacks the migration protein by administering an exogenous polynucleotide that encodes a migration protein. In a target that contains a migration protein, one can also identify administered agents that increase or decrease the migration of a developing gonadal cell. One can also treat the genetic material of the target organism using standard methods and treatments and can then identify genetic changes that increase or decrease migration of developing gonadal cells.

CLAIMS

WE CLAIM:

1. A method for identifying a modulator of a protein that comprises a metalloprotease domain and a
5 thrombospondin domain, the method comprising the steps of:
treating a target organism having a developing gonadal cell responsive to the protein with at least one potential modulator of cell migration; and
observing in the treated target organism a change in
10 migration or shape of the developing gonadal cell
attributable to the presence of the at least one modulator.
2. A method as claimed in Claim 1 wherein migration of the developing gonadal cell in the target organism before treatment is absent or reduced relative to a wild
15 type individual.
3. A method as claimed in Claim 1 wherein the treating step restores or enhances migration in the target organism relative to migration before the treating step.
4. A method as claimed in Claim 1 wherein migration
20 of the developing gonadal cell in the target organism before treatment is at a level of a wild type individual.
5. A method as claimed in Claim 1 wherein the treating step reduces migration in the target organism relative to migration before the treating step.

6. A method as claimed in Claim 1 wherein the target organism comprises a protein that comprises a metalloprotease domain and a thrombospondin domain, the protein being selected from the group consisting of a
5 protein encoded by a native polynucleotide coding sequence, a protein encoded by a heterologous polynucleotide coding sequence introduced into the target organism, a protein that shares at least 20% amino acid sequence identity with either of the foregoing and retains an ability to direct
10 cell migration in the target organism, and a chimeric protein encoded at least in part by at least one of the foregoing and introduced into the target organism, the polynucleotide coding sequence being under transcriptional control of a promoter active in a tissue located
15 sufficiently close to the developing gonadal cell so as to signal the cell to migrate.

7. A method as claimed in Claim 6, wherein the native polynucleotide coding sequence is *C. elegans gon-1*.

8. A method as claimed in Claim 6, wherein the
20 heterologous polynucleotide coding sequence is a homolog of *C. elegans gon-1*.

9. A method as claimed in Claim 8 wherein the homolog of *C. elegans gon-1* encodes a metalloprotease enzyme selected from the group consisting of murine ADAMTS-1
25 protein, bovine procollagen-1 N-proteinase, and human aggrecan-degrading metalloprotease.

10. A method as claimed in Claim 6 wherein the protein is truncated relative to a protein in a wild type individual.

11. A method as claimed in Claim 1 wherein the target organism is a nematode.

12. A method as claimed in Claim 11 wherein the target organism is a nematode selected from the group consisting
5 of *C. elegans* and *C. briggsae*.

13. A method as claimed in Claim 1 wherein the at least one modulator is selected from the group consisting of a nucleic acid molecule, a protein molecule, a sugar, a lipid, an organic molecule, a synthetic or natural
10 pharmaceutical agent, and a mixture thereof.

14. A method for identifying a nucleic acid sequence that affects migration of a developing gonadal cell, the method comprising the steps of:

treating a target organism by a method selected from
15 the group consisting of RNA interference, reverse genetics, and chemical mutagenesis to alter migration or shape of the developing gonadal cell in the treated target organism relative to migration in the target organism before treatment; and

20 identifying in the treated target organism a nucleic acid sequence affected by the treating step.

15. A method as claimed in Claim 14 wherein the treating step affects a nucleic acid sequence that encodes a protein.

16. A method as claimed in Claim 14 wherein the treating step affects a nucleic acid sequence that regulates nucleic acid transcription or translation.

17. A method as claimed in Claim 14 wherein migration
5 of the developing gonadal cell in the target organism before treatment is absent or reduced relative to a wild type individual.

18. A method as claimed in Claim 14 wherein the
10 treating step restores or enhances migration of the developing gonadal cell in the treated target organism relative to migration before the treating step.

19. A method as claimed in Claim 14 wherein migration of the developing gonadal cell in the target organism before treatment is at a level of a wild type individual.

15 20. A method as claimed in Claim 14 wherein the treating step reduces migration of the developing gonadal cell in the treated target organism relative to migration before the treating step.

21. A method as claimed in Claim 14, wherein the target organism comprises a protein that directs cell migration, the protein being selected from the group consisting of a protein encoded by a native polynucleotide coding sequence, a protein encoded by a heterologous polynucleotide coding sequence introduced into the target organism, a protein that shares at least 20% amino acid sequence identity with either of the foregoing and retains an ability to direct cell migration in the target organism, and a chimeric protein encoded at least in part by at least one of the foregoing and introduced into the target organism, the polynucleotide coding sequence being under transcriptional control of a promoter active in a tissue located sufficiently close to the developing gonadal cell so as to signal the cell to migrate.

22. A method as claimed in Claim 21 wherein the native polynucleotide coding sequence is *C. elegans gon-1*.

23. A method as claimed in Claim 21 wherein the heterologous polynucleotide coding sequence is a homolog of *C. elegans gon-1*.

24. A method as claimed in Claim 23 wherein the homolog of *C. elegans gon-1* encodes a metalloprotease enzyme selected from the group consisting of murine ADAMTS-1 protein, bovine procollagen-1 N-proteinase, and human aggrecan-degrading metalloprotease.

25. A method as claimed in Claim 21 wherein the protein is truncated relative to a protein in the wild type individual.

26. A method as claimed in Claim 14 wherein the target organism is a nematode.

27. A method as claimed in Claim 26 wherein the target organism is a nematode selected from the group consisting
5 of *C. elegans* and *C. briggsae*.

Fig. 1A




Fig. 1B Domains: MP TSpt1 TSpt1-like





Fig. 1C

GON-1 ADAMTS-1 PINP	signal sequence		119 122 83
	[F]R[S]G[G]F[H]L[L]Q[P]V[V]A[L]L[L]V[L]C[L]V[Y]A[L]Q[S]E[G]T[I]S[E]S[D]V[L]F[S]R[A]K[Y]G[V]P[V]H[S]R[V]Q[D]A[G]I[H]T[D]S[H]I[V]R[R]D[S]Y[G]R[R]G[K]R.....D[V]F[T]S[D]R[R]R[L]O[G]V[A]R[D]C[H]A[H]R[L]R[S]D[D]A[V]Y		
	[M]G[D]V[R]A[R]S[R]G[S]L[S]A[L]L[L]U[A]S[I]T[L]L[C]A[R]G[A].....H[G]R[P]T[E]E[D]E[L]M.....P[E]S[L]E[R]A[P]G[H].....D[S]H[T]T.....R[L]R[L]D[A]F.....[E]Q[Q]H[L]K[L]P[D]S[E]F		
GON-1 ADAMTS-1 PINP	[M]D[P]P[A]G[A]G[R]L[L]C[P]A[L]L[L]L[L]L[L]P[L]P[A]D[A]R[L]A[A]A[A]D[P]P[G]G[P]O[G]H[G]A[E]R[L]A[V]P[V]R[T]D[A]G[R]L[V]S[H]V[S]A[T]A[P]A[G]V[R]Q[R]R[A]A[P]A[Q]I[P]G[L]S[G]G[E]E[D]P[G]G[R]L[F]Y[N]V[T]V[E]F[R]D[L]H[R]L[R]P[N]A[R]L		
	[I]V[H]L[H]R[N]Q[I]P[D]S[L]N[K]S[V]P[H]F[S]N[S]N[F]A[P]V[L]L[D]S[E]E[V]R[G]G[F]R[T]D[P]D[L]I[Y]R[A]H[V]K[G]V[H]D[S]I[N]L[C]S[E]D[G]L[Y]G[M]L[P]S[G]H[T]V[E]P[I]T[S]G[N]G[T]E[D]G[A]S[R]H[R]Q[H]L[V]R[K]F[D]P[M]H[F]K[S]F[D]H[L]		
	[L]A[P]G[F]T[L]Q[V]G[R]P[G]S[E]A[Q]H[L]D[P]T[G]D[L]A[N]C[F]P[G]G[T]V[N]G[D]P[S]A[L]A[L]S[L]E[G]V[R]G[A]F[Y]L[Q]S[E]E[F]F[T]Q[P]A[G]V[A]T[E]R[L]A[P]A[P]E[E]S[S]A[R]P[O]F[H]I[L]R[R]R[R]G[S]G[A]K[G]C[G]V[N]D[D]E[L]T[L]P[T]S[D]S[R]P[E]		
GON-1 ADAMTS-1 PINP	[V]A[P]G[A]T[V]E[N]Q[G]E[G]A[T]R[V]E[L].....L[G]T[C]L[V]G[D]V[A]G[L]A[E]S[S]V[A]L[S]N[C]D[G]L[A]G[L]I[R]M[E]E[E]F[F]E[L].....E[K]G[L]A[K]E[A]E[Q]R[V]H[V]V[H]R[P]T[S]D[P]P[L]G[G]P[Q]A[L].....D[T]G[I]S[A]D[S]L[D]S		
	[N]S[T]S[V]N[E]T[E]T[T]V[A]T[H]O[D]Q[Q]E[D]V[I]E[R]K[A]R[S]A[N]S[M]D[H]I[V]E[L]V[A]D[I]T[V]E[H]G[S]R[S].....L[E]D[I]M[L]T[L]F[S]T[A]S[I]P[H]Q[S]L[R]A[S]I[N]M[M]K[L]I[V]L[K]T[E]N[A]G[P]R[L]TQ[N]A[Q]I[L]Q[F]C[H]A[H]P[Q]Y[Y]N[D]F		
	[S]O[N]T[R]N[Q]P[V]R[D]P[T]Q[D]A[G]K[P]S[G]P[S]I[R]K[D]F[V]S[S]P[R]I[V]E[H]I[V]A[D]S[M]A[D]H[G]S[G]L[K]H[L]L[L]T[L]F[S]V[A]R[F]K[H]P[S]I[R]N[S]I[S]I[M]M[K]I[L]V[I]T[E]E[Q]K[G]P[E]M[T]S[A]A[L]I[L]E[N]V[F]C[S]M[K]Q[H]M[S]F		
GON-1 ADAMTS-1 PINP	[L]S[R]A[L]G[V]L[E].....E[R]V[N]S[R]R[R]R[R]R[R]A[A]D[D]Y[N]I[E]M[L]L[G]V[D]D[S]V[V]Q[F]H[G]T[E]H[V]Q[K]L[L]T[L]M[I]N[I]V[E]I[H]D[E]S[L]G[A]H[E]N[V]M[L]R[I]I[L]L[S]Y[G]K[S]M[S]L[E]I[G]N[S]O[S]L[E]N[V]C[R]M[A]Y[L]Q[D]K		
	[D]D[S]S[V]D[H]H[V]A[L]L[L]R[C]I[R]S[Q]G[K]C[D]T[L]E[L]A[E]L[G]T[M]D[M]Q[K]S[A]T[I]E[D]N[G]L[S]A[F]T[I]A[H]E[L]G[H]V[F]S[I]P[H]D[E]R[K]C[S]T[Y]M[P]V[N]K[N]F[H]I[M]A[P]T[E]Y[N]T[P]W[S]N[S]P[C]S[A]G[M]L[E]R[F]E[N]N[R]G[Q]T[Q]		
	[S]O[R]D[P]E[H]Y[D]I[A]L[L]F[R]O[D]L[C]G[S]H[T]C[D]T[L]E[M]A[D]V[G]T[V]C[D]P[S]R[S]O[S]W[I]E[D]D[G]L[Q]A[F]T[T]A[H]E[L]G[H]V[F]N[P]H[D]D[A]K[H]C[A]S[L]N[G]V[S]D[S]H[L]M[A]S[L]S[L]D[S]Q[P]A[S]P[C]S[A]Y[M]V[T]S[F]L[D]N[G]H[G].....E[D]		
GON-1 ADAMTS-1 PINP	[D]T[D]H[D]E[Y]H[L]A[L]F[L]T[R]O[F].....G[P]S[G]M[Q]E[A]P[AT]G[M]C[H]P[V]R[S]Q[L]N[H]E[D]G[F]S[A]F[V]V[A]H[E]T[G]H[V]L[G]M[E]H[D]G[Q].....G[N]R[C]G[D]E[V]R[L]G[S]I[M]A[P]L[Y]Q[A]A[F]R[F]H[S]K[E]S[Q]E[L]S[R]V[L]H[S].....Y[D]Q		
	[L]E[D]F[V]E[R]R[Y]E[D]V[F]R[D]E[F]K[K]D[A]H[Q]Q[K]F[F]E[P]A[S]E[L]D[P]V[M]P.....T[R]R[L]R[Q]A[T]F[Y]G[S]O[N]E[R]T[I]Q[P]M[A]D[G]T[P]D[E]S[R]M[H]F[H]E[A]C[H]V[R]L[A].....P[E]S[L]T[K]I[D]D[E]M[D]A[R]S[N]E[C]S[R]T[C]G[E]		
	[L]M[A]K[E]O[N].....P[I]K[L]P[S]O[L]P[E]T[L]D[A]N[R]Q[D]T[F]E[E]S[K]H[P]D[A]S[L]T[T]L[W]G[T]S[G]L[L]V[D]T[Q]H[F]P[A]D[G]I[S]G[E]G[L].....M[V]S[E]K[V]N[K]T[M]K[H]F[A]T[P]V[H]S[S]I[G]M[P]G[P]M[E]D[C]S[R]T[C]G[E]		
GON-1 ADAMTS-1 PINP	[L]R[E]D[F]T[H]O[M]P[A].....L[P]Q[L]P[E]L[H]S[M]N[E]D[R]E[F]E[G]Y[M]M[A]T[A]R[T]F[D]P[K]Q[L]W[O]S[H]P[D]N[P]Y[F].....D[K]T[G]S[G]P[L]D[G]T[C]A[P]E[H]H[L]F[K]E[C]H[L]T[P].....D[I]L[K]R[D]E[V]M[A]M[S]P[H]E[S]C[S]R[T]C[G]E		
	[P]O[K]G[L]R[C]I[S]P[K]Y[N]G[E]K[V]M[G]O[R]E[R]R[S]C[N]T[R]E[E]P[N]O[T]Q[P]Y[R]E[N]Q[S]E[F]N[K]D[G]I[G]Q[V]A[S]T[N]H[V]P[K]Y[A]N[V]A[P]N[E]R[C]K[L]V[R]L[S]G[S]A[F]Y[L]E[R]D[K]M[D]G[T]P[D].....R[N]G[D]D[L]E[V]A[E]A[L]M[P]A[E]		
	[M]O[Y]T[R]E[L]D[N]M[P]N[G]G[K]V[E]K[R]V[R]S[C]N[T]E[D]P[N]G[K]T[F]R[E]E[D]C[E]A[N]E[F]S[K]A[S]F[G]N[E]P[T]V[E]M[P]K[Y]A[G]Y[S]P[K]D[R]C[K]L[L]E[A]K[G]I[G]N[F]F[L]Q[P]M[D]G[T]P[D]S[P]O[S]T[S]M[E]V[E]D[E]D[K]K[A]G		
GON-1 ADAMTS-1 PINP	[K]F[R]T[L]E[N]P[P]A[N]G[G]T[C]S[L]A[Y]D[F]O[L]E[N]S[D]E[F]I[D]A[L]O[F]R[E]E[D]C[R]O[N]D[L]Y.....F[E]H[G]D[A]Q[H]H[L]P[HE]H[R]D[A]K[E]R[C]H[L]M[E]S[K]E[T]E[V]V[S]K[R]M[A]D[G]T[R]S[Y]K[D]A[F]S[L]E[V]R[E]D[K]K[V]E		
	[C]D[H]Q[L]E[L]R[R]K[C]G[V]C[G]D[S]S[E]R[V]M[A]Q[I]F[N]E[Q]G[T]F[S]Y[N]E[M]K[I]P[A]S[A]M[D]C[R]O[K]G[Y]N[N]K[E]D[D]N[V]L[S]R[A]A[M]E[F]L[L]N[G]H[F]Q[V]S[L]A[R]Q[I]A[F]Q[D]T[V]L[E]T[S]G[S]D[A]I[T]E[R]U[N]G[T]E[P]T[R]S[D]I[Y]		
	[C]D[R]I[D]E[K]K[K]F[K]C[G]V[G]C[G]N[S]T[E]K[K]S[E]I[V]T[S]T[R]P[G]Y[H]D[V]T[P]A[G]T[N]E[V]K[H]R[N]Q[R]S[R]N[G]S[F]L[A]I[R]A[A]D[E]T[Y]I[L]N[G]N[F]T[L]S[T]L[E]Q[D]L[T]Y[K]G[V]L[R]Y[S]G[S]S[A]A[L]E[R]I[T]R[S]F[S]L[K]E[P]I[T]		
GON-1 ADAMTS-1 PINP	[C]D[E]V[G]E[S]K[Q]E[K]C[G]V[G]C[G]D[N]S[E]K[V]M[A]Q[I]F[S]R[S]P[K]K[L]G[Y]K[F]E[L]P[A]G[A]R[H]L[L]I[D]E[A]D.....T[T]S[H]L[A]V[K]N[L]E[T]E[K]F[I]N[E]E[N]D[V]P[N]S[K]T[F]I[A]N[G]V[E]M[E]Y[R]O[E]D[E]R[E]E[N]D[T]M[E]R[L]H[G]T[T]		
	[V]H[T]S[V]E[L]S[H]P[P]D[S]M[E]T[A]A[V]P[N]A[V]I[R]P[S]S.....A[L]Y[L]R[V]I[D]T[M]E[D]D[R]A[L]R[E]Q[S]Q[K]L[M]D[M]S[T]R[Q]S[H]D[R]N[C]D[N]V[L]K[K]Q[A]T[R]M[C]N[I]D[C]S[T]R[M]I[T]E[D]V[S]S[C]A[K]E[G]S[E]R[K]Q[C]		
	[I]D[M]N[V]G[A]L[R]P[K]I[K]F[H]F[M]K[K]T[E]S[F]...N[A]I...P[T]F[S]E[L]V[E]E[D]S[E]K[S]I[D]E[S]M[A]D[R]R[V]V[Q]D[D]I[N]G[P]A[S].....E[D]A[K]E[V]K[P]A[S]T[R]P[L]A[D]L[P]C[P]P[A]M[V]G[D]H[S]P[C]S[K]T[E]G[K]E[Y]K[K]R[C]		
GON-1 ADAMTS-1 PINP	[V]M[P]E[S]D[A]R[S]I[S]I[M]K[H]I[H]E[D]S[L]N[V]D[D]N[V]L[D]E[D]S[V]G[Y]E[L]A[L]K[K]A[S]P[E]S[K]P[E]G[S]D[F]T[K]Y[G]C[H]R[R]L[D]K[M]V[H]R[G]F[D]S[V]S[K]P[K]A[D]R[R]T[P]N[P]Q[E]S[Q]P[V]M[V]T[G]E[N]E[P]C[S]R[S]E[G]T[E]M[Q]V[R]		
	[R]W[S]V[K]H[E]G[D]R[Q]T[P]A[S]E[H]L[D]R[N]S[G]S[D]I[A]S[Q]Y[I]D[D].....plus 1148 aa		
	[T]L[K]C[V]S[H]D[G]G.....V[L]S[N]E[S]D[P]L[K]K[H]Y[D]F[L]T[L]O[C].....plus 1 aa		
	[S]V[R]C[V]P[L]H[N]T[T]R[S]V[H]T[K]H[N]D[A]R[E]G[R]R[A]G[N]R[E]L[D].....plus 236 aa		



SEQUENCE LISTING

<110> Kimble, Judith E
 Blelloch, Robert H

<120> Agent and Method for Modulating Cell Migration

5 <130> 960296.95386

<140>
 <141>

<150> 60/087170
 <151> 1998-05-29

10 <150> 60/129023
 <151> 1999-04-13

<160> 2

<170> PatentIn Ver. 2.0

15 <210> 1
 <211> 6659
 <212> DNA
 <213> Caenorhabditis elegans

20 <220>
 <221> CDS
 <222> (1)..(6453)

<400> 1
 atg cgc tcc atc ggc ggc tca ttc cat ctg ctg cag ccc gtc gtc gcc 48
 Met Arg Ser Ile Gly Gly Ser Phe His Leu Leu Gln Pro Val Val Ala
 1 5 10 15

25 gct ctc ata ctc ctc gtc gtc tgc ctc gtt tat gcg ttg caa tca ggg 96
 Ala Leu Ile Leu Leu Val Val Cys Leu Val Tyr Ala Leu Gln Ser Gly
 20 25 30

30 agt ggc acg atc tca gaa ttc tca tca gat gtg ctg ttc tcc agg gcc 144
 Ser Gly Thr Ile Ser Glu Phe Ser Ser Asp Val Leu Phe Ser Arg Ala
 35 40 45

	aag tac tca ggt gtg cca gtg cat cac agt cga tgg cgt caa gac gcc	192
	Lys Tyr Ser Gly Val Pro Val His His Ser Arg Trp Arg Gln Asp Ala	
	50 55 60	
5	ggt ata cac gtc atc gac agc cat cac atc gtc cga aga gat tct tat	240
	Gly Ile His Val Ile Asp Ser His His Ile Val Arg Arg Asp Ser Tyr	
	65 70 75 80	
	gga cgt cgt gga aaa cgt gat gtc acg tca aca gat cgg cga cgt cga	288
	Gly Arg Arg Gly Lys Arg Asp Val Thr Ser Thr Asp Arg Arg Arg Arg	
	85 90 95	
10	ctc caa gga gtt gcc aga gac tgt gga cat gct tgt cac tta cga tta	336
	Leu Gln Gly Val Ala Arg Asp Cys Gly His Ala Cys His Leu Arg Leu	
	100 105 110	
15	cga tca gat gat gcc gtc tac atc gtt cat ttg cac aga tgg aat caa	384
	Arg Ser Asp Asp Ala Val Tyr Ile Val His Leu His Arg Trp Asn Gln	
	115 120 125	
	ata ccg gac tca cat aac aaa agt gtt ccc cac ttt tcc aat tca aat	432
	Ile Pro Asp Ser His Asn Lys Ser Val Pro His Phe Ser Asn Ser Asn	
	130 135 140	
20	ttc gcg ccg atg gtc tta tat ttg gac tcg gag gag gag gtt aga ggt	480
	Phe Ala Pro Met Val Leu Tyr Leu Asp Ser Glu Glu Glu Val Arg Gly	
	145 150 155 160	
	gga atg tct cga aca gat ccc gat tgt atc tac cgt gca cac gtt aaa	528
	Gly Met Ser Arg Thr Asp Pro Asp Cys Ile Tyr Arg Ala His Val Lys	
	165 170 175	
25	ggt gta cat cag cac agc atc gtc aat tta tgc gac tcg gaa gac gga	576
	Gly Val His Gln His Ser Ile Val Asn Leu Cys Asp Ser Glu Asp Gly	
	180 185 190	
30	ttg tac gga atg ctt gca cta ccc agc gga atc cat acg gtt gag cca	624
	Leu Tyr Gly Met Leu Ala Leu Pro Ser Gly Ile His Thr Val Glu Pro	
	195 200 205	
	att att agt gga aac gga aca gag cac gac gga gca agt cgc cat agg	672
	Ile Ile Ser Gly Asn Gly Thr Glu His Asp Gly Ala Ser Arg His Arg	
	210 215 220	

	caa cat ctc gtc cga aag ttc gat cca atg cac ttc aaa tcg ttt gac	720
	Gln His Leu Val Arg Lys Phe Asp Pro Met His Phe Lys Ser Phe Asp	
	225 230 235 240	
5	cat ctt aac tcg acc agt gtc aac gag acg gag acg acg gtt gcc acg	768
	His Leu Asn Ser Thr Ser Val Asn Glu Thr Glu Thr Thr Val Ala Thr	
	245 250 255	
	tgg caa gat cag tgg gaa gat gtt att gaa cgc aaa gca aga tcc cga	816
	Trp Gln Asp Gln Trp Glu Asp Val Ile Glu Arg Lys Ala Arg Ser Arg	
	260 265 270	
10	aga gct gcc aac tct tgg gat cac tat gtt gaa gtc ctt gtg gtg gcg	864
	Arg Ala Ala Asn Ser Trp Asp His Tyr Val Glu Val Leu Val Val Ala	
	275 280 285	
15	gat aca aaa atg tac gaa tat cac gga aga tct ctt gaa gac tac gtt	912
	Asp Thr Lys Met Tyr Glu Tyr His Gly Arg Ser Leu Glu Asp Tyr Val	
	290 295 300	
	ctc act ctc ttc tcc aca gtt gcc tcc atc tat cgt cac caa tcc ctt	960
	Leu Thr Leu Phe Ser Thr Val Ala Ser Ile Tyr Arg His Gln Ser Leu	
	305 310 315 320	
20	cgt gca tct atc aat gtc gtt gtt gtc aag ttg atc gtt ttg aaa acg	1008
	Arg Ala Ser Ile Asn Val Val Val Val Lys Leu Ile Val Leu Lys Thr	
	325 330 335	
	gaa aac gct gga cca cga atc act cag aac gct caa caa aca ctt caa	1056
	Glu Asn Ala Gly Pro Arg Ile Thr Gln Asn Ala Gln Gln Thr Leu Gln	
	340 345 350	
25	gat ttc tgt aga tgg cag cag tat tac aat gat cca gat gat tcg agt	1104
	Asp Phe Cys Arg Trp Gln Gln Tyr Tyr Asn Asp Pro Asp Asp Ser Ser	
	355 360 365	
30	gtc caa cat cat gac gtt gca atc ctt ttg acg cgt aaa gat att tgt	1152
	Val Gln His His Asp Val Ala Ile Leu Leu Thr Arg Lys Asp Ile Cys	
	370 375 380	
	cga tca caa gga aaa tgc gat aca ctt gga ctt gct gaa ctt gga aca	1200
	Arg Ser Gln Gly Lys Cys Asp Thr Leu Gly Leu Ala Glu Leu Gly Thr	
	385 390 395 400	

	atg tgt gat atg caa aaa agt tgt gca atc ata gaa gac aat gga ttg	1248
	Met Cys Asp Met Gln Lys Ser Cys Ala Ile Ile Glu Asp Asn Gly Leu	
	405 410 415	
5	agt gct gca ttc aca att gct cat gaa ttg ggt cat gtg ttt tcg att	1296
	Ser Ala Ala Phe Thr Ile Ala His Glu Leu Gly His Val Phe Ser Ile	
	420 425 430	
	cct cat gat gac gaa cga aaa tgc tct acc tac atg ccg gtt aat aag	1344
	Pro His Asp Asp Glu Arg Lys Cys Ser Thr Tyr Met Pro Val Asn Lys	
	435 440 445	
10	aac aac ttc cac ata atg gca cca acg ttg gaa tat aac act cat cca	1392
	Asn Asn Phe His Ile Met Ala Pro Thr Leu Glu Tyr Asn Thr His Pro	
	450 455 460	
15	tgg agt tgg tcg cca tgt tca gct gga atg ctc gaa cga ttc ctc gaa	1440
	Trp Ser Trp Ser Pro Cys Ser Ala Gly Met Leu Glu Arg Phe Leu Glu	
	465 470 475 480	
	aat aat cga ggt caa act caa tgt cta ttc gat cag ccg gtc gaa cgt	1488
	Asn Asn Arg Gly Gln Thr Gln Cys Leu Phe Asp Gln Pro Val Glu Arg	
	485 490 495	
20	cgt tac tac gag gat gtc ttt gta cgt gat gaa cca gga aag aaa tac	1536
	Arg Tyr Tyr Glu Asp Val Phe Val Arg Asp Glu Pro Gly Lys Lys Tyr	
	500 505 510	
	gat gct cat caa cag tgc aag ttt gta ttt gga cca gct tct gag ttg	1584
	Asp Ala His Gln Gln Cys Lys Phe Val Phe Gly Pro Ala Ser Glu Leu	
	515 520 525	
25	tgc cct tat atg ccg aca tgc cgc cgt ctt tgg tgt gca aca ttc tac	1632
	Cys Pro Tyr Met Pro Thr Cys Arg Arg Leu Trp Cys Ala Thr Phe Tyr	
	530 535 540	
30	gga agc cag atg ggc tgt cga act cag cat atg cca tgg gcc gac gga	1680
	Gly Ser Gln Met Gly Cys Arg Thr Gln His Met Pro Trp Ala Asp Gly	
	545 550 555 560	
	act cct tgt gac gaa tca aga agc atg ttc tgt cat cat gga gcc tgt	1728
	Thr Pro Cys Asp Glu Ser Arg Ser Met Phe Cys His His Gly Ala Cys	
	565 570 575	

	gtt cgt cta gcc ccc gaa tcc ctt acc aaa att gac gga caa tgg ggt	1776
	Val Arg Leu Ala Pro Glu Ser Leu Thr Lys Ile Asp Gly Gln Trp Gly	
	580 585 590	
5	gac tgg cga tca tgg gga gaa tgc agt cgt act tgt ggt ggt ggt gtt	1824
	Asp Trp Arg Ser Trp Gly Glu Cys Ser Arg Thr Cys Gly Gly Gly Val	
	595 600 605	
	caa aaa gga tta aga gat tgt gac agc cca aaa cct cga aat ggt gga	1872
	Gln Lys Gly Leu Arg Asp Cys Asp Ser Pro Lys Pro Arg Asn Gly Gly	
	610 615 620	
10	aag tac tgt gtt ggt caa cga gaa cgt tat cgg tca tgt aat aca caa	1920
	Lys Tyr Cys Val Gly Gln Arg Glu Arg Tyr Arg Ser Cys Asn Thr Gln	
	625 630 635 640	
15	gaa tgc cca tgg gat act caa cca tac cgt gaa gtt caa tgt tct gaa	1968
	Glu Cys Pro Trp Asp Thr Gln Pro Tyr Arg Glu Val Gln Cys Ser Glu	
	645 650 655	
	ttc aac aat aaa gat att gga atc caa ggt gtc gct tca acg aat act	2016
	Phe Asn Asn Lys Asp Ile Gly Ile Gln Gly Val Ala Ser Thr Asn Thr	
	660 665 670	
20	cac tgg gtt cca aaa tat gcg aat gtt gca cca aat gaa cgt tgc aag	2064
	His Trp Val Pro Lys Tyr Ala Asn Val Ala Pro Asn Glu Arg Cys Lys	
	675 680 685	
	ctg tat tgt cgg ctc agt gga tct gca gcg ttc tat ctg ctt cga gat	2112
	Leu Tyr Cys Arg Leu Ser Gly Ser Ala Ala Phe Tyr Leu Leu Arg Asp	
	690 695 700	
25	aaa gtt gtt gat gga aca cca tgt gat aga aat gga gac gat att tgt	2160
	Lys Val Val Asp Gly Thr Pro Cys Asp Arg Asn Gly Asp Asp Ile Cys	
	705 710 715 720	
30	gta gct gga gct tgt atg cca gca ggc tgt gat cat caa ctt cat tca	2208
	Val Ala Gly Ala Cys Met Pro Ala Gly Cys Asp His Gln Leu His Ser	
	725 730 735	
	act ctc cga aga gac aaa tgt ggt gtt tgc ggt ggg gat gat tct tcc	2256
	Thr Leu Arg Arg Asp Lys Cys Gly Val Cys Gly Gly Asp Asp Ser Ser	
	740 745 750	

	tgt aag gtt gtc aaa gga aca ttt aat gag caa gga acc ttt ggt tat	2304
	Cys Lys Val Val Lys Gly Thr Phe Asn Glu Gln Gly Thr Phe Gly Tyr	
	755 760 765	
5	aac gaa gta atg aag att cca gct ggt tct gca aat att gat atc cgg	2352
	Asn Glu Val Met Lys Ile Pro Ala Gly Ser Ala Asn Ile Asp Ile Arg	
	770 775 780	
	cag aaa gga tat aat aat atg aaa gaa gat gac aat tat ctt tct ctc	2400
	Gln Lys Gly Tyr Asn Asn Met Lys Glu Asp Asp Asn Tyr Leu Ser Leu	
	785 790 795 800	
10	cgt gcc gcc aat ggt gaa ttc cta ctt aac ggt cat ttc caa gta tca	2448
	Arg Ala Ala Asn Gly Glu Phe Leu Leu Asn Gly His Phe Gln Val Ser	
	805 810 815	
15	ctg gct cgc caa caa att gca ttc caa gac act gtt ctc gaa tat tct	2496
	Leu Ala Arg Gln Gln Ile Ala Phe Gln Asp Thr Val Leu Glu Tyr Ser	
	820 825 830	
	ggt tct gat gca att att gaa cgg ata aat gga act ggt ccg att aga	2544
	Gly Ser Asp Ala Ile Ile Glu Arg Ile Asn Gly Thr Gly Pro Ile Arg	
	835 840 845	
20	agt gac att tat gtt cat gtt ctt tct gtt ggt agt cat cca ccc gac	2592
	Ser Asp Ile Tyr Val His Val Leu Ser Val Gly Ser His Pro Pro Asp	
	850 855 860	
	atc tca tat gag tac atg act gcg gct gtt cca aat gct gta att cgg	2640
	Ile Ser Tyr Glu Tyr Met Thr Ala Ala Val Pro Asn Ala Val Ile Arg	
	865 870 875 880	
25	cca ata tcc agt gca ttg tat ttg tgg aga gtt acg gat act tgg aca	2688
	Pro Ile Ser Ser Ala Leu Tyr Leu Trp Arg Val Thr Asp Thr Trp Thr	
	885 890 895	
30	gaa tgt gat aga gcc tgt cgt gga cag caa tcg caa aaa tta atg tgt	2736
	Glu Cys Asp Arg Ala Cys Arg Gly Gln Gln Ser Gln Lys Leu Met Cys	
	900 905 910	
	ctg gac atg tcg act cat cgt caa agt cat gat aga aat tgt caa aat	2784
	Leu Asp Met Ser Thr His Arg Gln Ser His Asp Arg Asn Cys Gln Asn	
	915 920 925	
	gtt ctc aaa cca aaa caa gca aca cga atg tgc aat ata gat tgt tct	2832

	Val	Leu	Lys	Pro	Lys	Gln	Ala	Thr	Arg	Met	Cys	Asn	Ile	Asp	Cys	Ser	
	930						935					940					
5	aca	aga	tgg	atc	act	gaa	gat	gtg	tct	agt	tgt	agt	gcc	aaa	tgt	gga	2880
	Thr	Arg	Trp	Ile	Thr	Glu	Asp	Val	Ser	Ser	Cys	Ser	Ala	Lys	Cys	Gly	
	945					950					955					960	
	tct	gga	cag	aaa	cgt	caa	cga	gtt	tct	tgc	gta	aaa	atg	gag	ggt	gat	2928
	Ser	Gly	Gln	Lys	Arg	Gln	Arg	Val	Ser	Cys	Val	Lys	Met	Glu	Gly	Asp	
					965					970					975		
10	cgt	caa	act	cca	gca	tcc	gaa	cat	cta	tgt	gat	cgt	aat	tca	aaa	cca	2976
	Arg	Gln	Thr	Pro	Ala	Ser	Glu	His	Leu	Cys	Asp	Arg	Asn	Ser	Lys	Pro	
				980					985					990			
	tcc	gat	att	gcc	agt	tgt	tac	att	gac	tgc	tct	gga	aga	aaa	tgg	aac	3024
	Ser	Asp	Ile	Ala	Ser	Cys	Tyr	Ile	Asp	Cys	Ser	Gly	Arg	Lys	Trp	Asn	
			995				1000					1005					
15	tat	gga	gaa	tgg	act	tca	tgt	tct	gaa	act	tgc	gga	tcg	aat	gga	aaa	3072
	Tyr	Gly	Glu	Trp	Thr	Ser	Cys	Ser	Glu	Thr	Cys	Gly	Ser	Asn	Gly	Lys	
		1010					1015				1020						
	atg	cat	cgg	aag	tca	tat	tgc	gtt	gat	gat	tcg	aat	cgt	cga	gtt	gat	3120
20	Met	His	Arg	Lys	Ser	Tyr	Cys	Val	Asp	Asp	Ser	Asn	Arg	Arg	Val	Asp	
	1025					1030				1035					1040		
	gag	tca	ttg	tgc	ggc	aga	gaa	cag	aaa	gag	gcg	aca	gaa	cgg	gaa	tgt	3168
	Glu	Ser	Leu	Cys	Gly	Arg	Glu	Gln	Lys	Glu	Ala	Thr	Glu	Arg	Glu	Cys	
				1045					1050					1055			
25	aac	aga	att	cca	tgt	cca	aga	tgg	gtt	tat	ggg	cat	tgg	tca	gag	tgc	3216
	Asn	Arg	Ile	Pro	Cys	Pro	Arg	Trp	Val	Tyr	Gly	His	Trp	Ser	Glu	Cys	
				1060				1065					1070				
	tct	cga	agt	tgt	gat	ggt	gga	gtc	aaa	atg	cgt	cat	gct	caa	tgt	ttg	3264
	Ser	Arg	Ser	Cys	Asp	Gly	Gly	Val	Lys	Met	Arg	His	Ala	Gln	Cys	Leu	
			1075				1080					1085					
30	gat	gca	gcc	gat	cgg	gaa	aca	cat	aca	tcc	aga	tgt	ggt	cca	gca	cag	3312
	Asp	Ala	Ala	Asp	Arg	Glu	Thr	His	Thr	Ser	Arg	Cys	Gly	Pro	Ala	Gln	
		1090				1095					1100						

	aca caa gaa cat tgt aat gaa cat gct tgt act tgg tgg cag ttc gga	3360
	Thr Gln Glu His Cys Asn Glu His Ala Cys Thr Trp Trp Gln Phe Gly	
	1105 1110 1115 1120	
5	gtc tgg tct gac tgc tca gct aag tgt gga gat ggt gta cag tat cga	3408
	Val Trp Ser Asp Cys Ser Ala Lys Cys Gly Asp Gly Val Gln Tyr Arg	
	1125 1130 1135	
	gac gct aat tgt acc gat cgt cat aga tca gta cta ccg gaa cat cgt	3456
	Asp Ala Asn Cys Thr Asp Arg His Arg Ser Val Leu Pro Glu His Arg	
	1140 1145 1150	
10	tgc ctt aaa atg gaa aag ata att aca aaa cca tgt cat aga gaa tca	3504
	Cys Leu Lys Met Glu Lys Ile Ile Thr Lys Pro Cys His Arg Glu Ser	
	1155 1160 1165	
15	tgt cca aaa tat aaa ctt gga gaa tgg tct cag tgt agt gtt tct tgt	3552
	Cys Pro Lys Tyr Lys Leu Gly Glu Trp Ser Gln Cys Ser Val Ser Cys	
	1170 1175 1180	
	gag gat gga tgg tcg tca aga aga gtt tca tgt gtt tct gga aat gga	3600
	Glu Asp Gly Trp Ser Ser Arg Arg Val Ser Cys Val Ser Gly Asn Gly	
	1185 1190 1195 1200	
20	act gaa gtc gat atg tca ctt tgt ggt act gca tct gat cgg cct gct	3648
	Thr Glu Val Asp Met Ser Leu Cys Gly Thr Ala Ser Asp Arg Pro Ala	
	1205 1210 1215	
	tct cat cag aca tgt aat tta ggc act tgc cca ttt tgg aga aat act	3696
	Ser His Gln Thr Cys Asn Leu Gly Thr Cys Pro Phe Trp Arg Asn Thr	
	1220 1225 1230	
25	gat tgg agt gct tgt tct gta tct tgt gga atc ggt cat cgg gaa cgt	3744
	Asp Trp Ser Ala Cys Ser Val Ser Cys Gly Ile Gly His Arg Glu Arg	
	1235 1240 1245	
30	aca acc gaa tgc ata tac cgc gaa caa tct gtt gat gct tct ttt tgt	3792
	Thr Thr Glu Cys Ile Tyr Arg Glu Gln Ser Val Asp Ala Ser Phe Cys	
	1250 1255 1260	
	gga gat acc aaa atg cca gaa act agt caa act tgc cat ctt ctg cca	3840
	Gly Asp Thr Lys Met Pro Glu Thr Ser Gln Thr Cys His Leu Leu Pro	
	1265 1270 1275 1280	

	tgt aca tct tgg aaa cca agt cat tgg tcc cct tgc tca gtc act tgt	3888
	Cys Thr Ser Trp Lys Pro Ser His Trp Ser Pro Cys Ser Val Thr Cys	
	1285 1290 1295	
5	gga tca gga att cag act aga agt gtt tgc tgt act cgt gga tct gaa	3936
	Gly Ser Gly Ile Gln Thr Arg Ser Val Ser Cys Thr Arg Gly Ser Glu	
	1300 1305 1310	
	gga act att gtt gat gaa tat ttt tgt gat cga aat act cgt cca cgc	3984
	Gly Thr Ile Val Asp Glu Tyr Phe Cys Asp Arg Asn Thr Arg Pro Arg	
	1315 1320 1325	
10	cta aaa aag act tgt gaa aaa gat act tgt gat ggg ccc aga gta ctt	4032
	Leu Lys Lys Thr Cys Glu Lys Asp Thr Cys Asp Gly Pro Arg Val Leu	
	1330 1335 1340	
	caa aaa ctt caa gcc gac gta cca cca atc cga tgg gca acc gga cca	4080
15	Gln Lys Leu Gln Ala Asp Val Pro Pro Ile Arg Trp Ala Thr Gly Pro	
	1345 1350 1355 1360	
	tgg aca gcc tgt tca gca act tgt ggt aat ggt act caa cgt cgt ctt	4128
	Trp Thr Ala Cys Ser Ala Thr Cys Gly Asn Gly Thr Gln Arg Arg Leu	
	1365 1370 1375	
20	ctc aag tgc cga gat cat gtt cgt gat ctt cct gat gag tat tgc aat	4176
	Leu Lys Cys Arg Asp His Val Arg Asp Leu Pro Asp Glu Tyr Cys Asn	
	1380 1385 1390	
	cat ttg gat aag gaa gta tca aca aga aat tgt cgc ctt cgt gat tgt	4224
	His Leu Asp Lys Glu Val Ser Thr Arg Asn Cys Arg Leu Arg Asp Cys	
	1395 1400 1405	
25	tca tac tgg aaa atg gcg gaa tgg gaa gag tgt cca gct act tgt gga	4272
	Ser Tyr Trp Lys Met Ala Glu Trp Glu Glu Cys Pro Ala Thr Cys Gly	
	1410 1415 1420	
	act cat gtt caa caa agt aga aat gtt aca tgc gtc agt gcg gaa gac	4320
30	Thr His Val Gln Gln Ser Arg Asn Val Thr Cys Val Ser Ala Glu Asp	
	1425 1430 1435 1440	
	ggt ggt cgg acg att ttg aaa gat gtt gat tgt gat gtg caa aag aga	4368
	Gly Gly Arg Thr Ile Leu Lys Asp Val Asp Cys Asp Val Gln Lys Arg	
	1445 1450 1455	

	cca	aca	agt	gca	aga	aat	tgc	cga	ctt	gaa	ccc	tgt	cca	aag	gga	gaa	4416
	Pro	Thr	Ser	Ala	Arg	Asn	Cys	Arg	Leu	Glu	Pro	Cys	Pro	Lys	Gly	Glu	
				1460					1465							1470	
5	gaa	cat	att	gga	tcc	tgg	att	att	gga	gat	tgg	tca	aaa	tgc	tct	gct	4464
	Glu	His	Ile	Gly	Ser	Trp	Ile	Ile	Gly	Asp	Trp	Ser	Lys	Cys	Ser	Ala	
				1475					1480							1485	
	tct	tgt	ggt	ggg	gga	tgg	cgt	cgt	cgc	agt	gta	tct	tgc	act	tcg	tct	4512
	Ser	Cys	Gly	Gly	Gly	Trp	Arg	Arg	Arg	Ser	Val	Ser	Cys	Thr	Ser	Ser	
				1490					1495							1500	
10	tct	tgc	gat	gaa	acc	aga	aaa	cca	aag	atg	ttt	gat	aaa	tgc	aat	gaa	4560
	Ser	Cys	Asp	Glu	Thr	Arg	Lys	Pro	Lys	Met	Phe	Asp	Lys	Cys	Asn	Glu	
				1505					1510							1520	
	gaa	cta	tgt	cca	cca	ctc	aca	aat	aat	tct	tgg	cag	ata	tct	cca	tgg	4608
15	Glu	Leu	Cys	Pro	Pro	Leu	Thr	Asn	Asn	Ser	Trp	Gln	Ile	Ser	Pro	Trp	
						1525					1530					1535	
	act	cac	tgt	tct	gta	tcg	tgt	ggc	ggg	gga	gtt	caa	cgc	cgc	aaa	atc	4656
	Thr	His	Cys	Ser	Val	Ser	Cys	Gly	Gly	Gly	Val	Gln	Arg	Arg	Lys	Ile	
						1540					1545					1550	
20	tgg	tgt	gaa	gac	gtg	ctt	tcc	ggt	cgt	aaa	caa	gac	gat	atc	gag	tgc	4704
	Trp	Cys	Glu	Asp	Val	Leu	Ser	Gly	Arg	Lys	Gln	Asp	Asp	Ile	Glu	Cys	
				1555					1560							1565	
	tca	gag	att	aag	cct	cgc	gaa	caa	aga	gat	tgt	gaa	atg	cct	cca	tgc	4752
	Ser	Glu	Ile	Lys	Pro	Arg	Glu	Gln	Arg	Asp	Cys	Glu	Met	Pro	Pro	Cys	
				1570					1575							1580	
25	cga	tct	cat	tat	cac	aac	aaa	aca	tca	tca	gca	tca	atg	aca	tca	tta	4800
	Arg	Ser	His	Tyr	His	Asn	Lys	Thr	Ser	Ser	Ala	Ser	Met	Thr	Ser	Leu	
						1585					1590					1600	
	tca	tct	tcg	aat	tca	aat	acg	acg	tct	tcc	gct	tcc	gct	tct	tcg	ctt	4848
30	Ser	Ser	Ser	Asn	Ser	Asn	Thr	Thr	Ser	Ser	Ala	Ser	Ala	Ser	Ser	Leu	
						1605					1610					1615	
	cct	atc	ctt	cca	ccc	gtc	gtc	tcc	tgg	caa	acg	tct	gca	tgg	agc	gcg	4896
	Pro	Ile	Leu	Pro	Pro	Val	Val	Ser	Trp	Gln	Thr	Ser	Ala	Trp	Ser	Ala	
						1620					1625					1630	

	tgt tct gca aaa tgc ggt cgt gga acg aaa cga aga gtt gtc gaa tgt	4944
	Cys Ser Ala Lys Cys Gly Arg Gly Thr Lys Arg Arg Val Val Glu Cys	
	1635 1640 1645	
5	gta aat cca tca tta aat gtg aca gtg gca agt aca gaa tgt gat caa	4992
	Val Asn Pro Ser Leu Asn Val Thr Val Ala Ser Thr Glu Cys Asp Gln	
	1650 1655 1660	
	acg aag aaa cca gtt gaa gaa gtt cgt tgt cgt act aaa cat tgc ccg	5040
	Thr Lys Lys Pro Val Glu Glu Val Arg Cys Arg Thr Lys His Cys Pro	
	1665 1670 1675 1680	
10	aga tgg aag act act act tgg agt tcg tgt tct gtc acc tgt ggc aga	5088
	Arg Trp Lys Thr Thr Thr Trp Ser Ser Cys Ser Val Thr Cys Gly Arg	
	1685 1690 1695	
15	gga atc aga cgt cgt gaa gtt caa tgt tat cgt ggt cgc aag aat ttg	5136
	Gly Ile Arg Arg Arg Glu Val Gln Cys Tyr Arg Gly Arg Lys Asn Leu	
	1700 1705 1710	
	gtg tct gat tcg gag tgc aat cca aaa act aag ctc aac tct gtt gcc	5184
	Val Ser Asp Ser Glu Cys Asn Pro Lys Thr Lys Leu Asn Ser Val Ala	
	1715 1720 1725	
20	aac tgt ttc cca gtg gct tgt cca gct tat aga tgg aat gtt act cca	5232
	Asn Cys Phe Pro Val Ala Cys Pro Ala Tyr Arg Trp Asn Val Thr Pro	
	1730 1735 1740	
	tgg agc aag tgc aaa gat gag tgt gct cga gga caa aag caa act cgt	5280
	Trp Ser Lys Cys Lys Asp Glu Cys Ala Arg Gly Gln Lys Gln Thr Arg	
	1745 1750 1755 1760	
25	cgg gtg cac tgt ata agc act tct ggt aaa cga gca gct cca cga atg	5328
	Arg Val His Cys Ile Ser Thr Ser Gly Lys Arg Ala Ala Pro Arg Met	
	1765 1770 1775	
30	tgt gaa ttg gct cgt gca cca act tcg atc aga gag tgc gat aca tca	5376
	Cys Glu Leu Ala Arg Ala Pro Thr Ser Ile Arg Glu Cys Asp Thr Ser	
	1780 1785 1790	
	aat tgt cca tat gag tgg gtg cca gga gat tgg caa acg tgt tca aag	5424
	Asn Cys Pro Tyr Glu Trp Val Pro Gly Asp Trp Gln Thr Cys Ser Lys	
	1795 1800 1805	

	tca tgt gga gaa gga gta cag aca cga gaa gtc aga tgt cgt aga aag	5472
	Ser Cys Gly Glu Gly Val Gln Thr Arg Glu Val Arg Cys Arg Arg Lys	
	1810 1815 1820	
5	att aat ttt aac tca acc att cca att ata ttt atg ctc gaa gat gaa	5520
	Ile Asn Phe Asn Ser Thr Ile Pro Ile Ile Phe Met Leu Glu Asp Glu	
	1825 1830 1835 1840	
	cca gct gta cca aaa gag aaa tgt gaa ctt ttc cca aaa cca aat gaa	5568
	Pro Ala Val Pro Lys Glu Lys Cys Glu Leu Phe Pro Lys Pro Asn Glu	
	1845 1850 1855	
10	tct caa acg tgc gaa ctt aac cca tgc gat tgc gaa ttc aaa tgg agt	5616
	Ser Gln Thr Cys Glu Leu Asn Pro Cys Asp Ser Glu Phe Lys Trp Ser	
	1860 1865 1870	
15	ttc gga cca tgg ggt gaa tgc tgc aaa aat tgc ggt caa ggt att cga	5664
	Phe Gly Pro Trp Gly Glu Cys Ser Lys Asn Cys Gly Gln Gly Ile Arg	
	1875 1880 1885	
	cgt cga cgt gtc aag tgt gtg gcc aat gat ggt cgt cga gtt gaa cga	5712
	Arg Arg Arg Val Lys Cys Val Ala Asn Asp Gly Arg Arg Val Glu Arg	
	1890 1895 1900	
20	gtc aag tgt acc aca aag aaa cca cgt cga act caa tat tgt ttt gaa	5760
	Val Lys Cys Thr Thr Lys Lys Pro Arg Arg Thr Gln Tyr Cys Phe Glu	
	1905 1910 1915 1920	
	aga aat tgc ctt ccg tca act tgt cag gag ctt aaa tct cag aat gtt	5808
	Arg Asn Cys Leu Pro Ser Thr Cys Gln Glu Leu Lys Ser Gln Asn Val	
	1925 1930 1935	
25	aag gct aaa gat gga aat tac act att ctt ctt gac gga ttc act att	5856
	Lys Ala Lys Asp Gly Asn Tyr Thr Ile Leu Leu Asp Gly Phe Thr Ile	
	1940 1945 1950	
30	gaa att tat tgt cat cga atg aat tca acc att cct aaa gct tat ttg	5904
	Glu Ile Tyr Cys His Arg Met Asn Ser Thr Ile Pro Lys Ala Tyr Leu	
	1955 1960 1965	
	aac gtt aat cca aga acc aat ttt gca gag gtt tat gga aaa aaa tta	5952
	Asn Val Asn Pro Arg Thr Asn Phe Ala Glu Val Tyr Gly Lys Lys Leu	
	1970 1975 1980	

	ata tac cct cat act tgc cca ttt aat ggt gat cgt aat gat tca tgc Ile Tyr Pro His Thr Cys Pro Phe Asn Gly Asp Arg Asn Asp Ser Cys 1985 1990 1995 2000	6000
5	cat tgt tca gaa gac ggc gat gca agt gct gga ttg acg aga ttc aat His Cys Ser Glu Asp Gly Asp Ala Ser Ala Gly Leu Thr Arg Phe Asn 2005 2010 2015	6048
	aaa gtt cga ata gat ttg ttg aat aga aag ttc cat ctg gcg gat tat Lys Val Arg Ile Asp Leu Leu Asn Arg Lys Phe His Leu Ala Asp Tyr 2020 2025 2030	6096
10	aca ttt gca aaa cga gaa tat ggt gtt cat gtg cca tat ggt act gcc Thr Phe Ala Lys Arg Glu Tyr Gly Val His Val Pro Tyr Gly Thr Ala 2035 2040 2045	6144
15	ggt gat tgc tac agt atg aaa gat tgt cca cag gga ata ttc tca att Gly Asp Cys Tyr Ser Met Lys Asp Cys Pro Gln Gly Ile Phe Ser Ile 2050 2055 2060	6192
	gat tta aaa tct gct ggt ctg aaa tta gtt gac gat ctg aat tgg gag Asp Leu Lys Ser Ala Gly Leu Lys Leu Val Asp Asp Leu Asn Trp Glu 2065 2070 2075 2080	6240
20	gat caa ggt cat cga aca tcc tct cga atc gat cgt ttt tat aac aat Asp Gln Gly His Arg Thr Ser Ser Arg Ile Asp Arg Phe Tyr Asn Asn 2085 2090 2095	6288
	gca aaa gtt att ggt cac tgt ggt ggt ttt tgt gga aaa tgc tct cct Ala Lys Val Ile Gly His Cys Gly Gly Phe Cys Gly Lys Cys Ser Pro 2100 2105 2110	6336
25	gag cgg tac aaa gga cta atc ttt gaa gtt aat aca aaa tta tta aat Glu Arg Tyr Lys Gly Leu Ile Phe Glu Val Asn Thr Lys Leu Leu Asn 2115 2120 2125	6384
30	cat gtg aaa aat ggt gga cac att gat gat gaa ttg gat gat gat ggt His Val Lys Asn Gly Gly His Ile Asp Asp Glu Leu Asp Asp Asp Gly 2130 2135 2140	6432
	ttc tct ggt gac atg gat taa ttttttcgat acctaaaagt gtcaaaatct Phe Ser Gly Asp Met Asp 2145 2150	6483

cgtatgaatc tctacttctc tggctctctta ttccaagttt ttgattcttt tctttttttt 6543
 agtttttaat agcattactt cgaatttatt gtcattccct caatcaccta acactagggt 6603
 ttctacatag tatgttcctt gaaaatgttt catgatcaaa ggttacggta cttttg 6659

5 <210> 2
 <211> 2150
 <212> PRT
 <213> Caenorhabditis elegans

10 <400> 2
 Met Arg Ser Ile Gly Gly Ser Phe His Leu Leu Gln Pro Val Val Ala
 1 5 10 15
 Ala Leu Ile Leu Leu Val Val Cys Leu Val Tyr Ala Leu Gln Ser Gly
 20 25 30
 Ser Gly Thr Ile Ser Glu Phe Ser Ser Asp Val Leu Phe Ser Arg Ala
 35 40 45
 15 Lys Tyr Ser Gly Val Pro Val His His Ser Arg Trp Arg Gln Asp Ala
 50 55 60
 Gly Ile His Val Ile Asp Ser His His Ile Val Arg Arg Asp Ser Tyr
 65 70 75 80
 20 Gly Arg Arg Gly Lys Arg Asp Val Thr Ser Thr Asp Arg Arg Arg Arg
 85 90 95
 Leu Gln Gly Val Ala Arg Asp Cys Gly His Ala Cys His Leu Arg Leu
 100 105 110
 Arg Ser Asp Asp Ala Val Tyr Ile Val His Leu His Arg Trp Asn Gln
 115 120 125
 25 Ile Pro Asp Ser His Asn Lys Ser Val Pro His Phe Ser Asn Ser Asn
 130 135 140
 Phe Ala Pro Met Val Leu Tyr Leu Asp Ser Glu Glu Glu Val Arg Gly
 145 150 155 160
 30 Gly Met Ser Arg Thr Asp Pro Asp Cys Ile Tyr Arg Ala His Val Lys
 165 170 175

	Gly Val His Gln His Ser Ile Val Asn Leu Cys Asp Ser Glu Asp Gly	180	185	190
	Leu Tyr Gly Met Leu Ala Leu Pro Ser Gly Ile His Thr Val Glu Pro	195	200	205
5	Ile Ile Ser Gly Asn Gly Thr Glu His Asp Gly Ala Ser Arg His Arg	210	215	220
	Gln His Leu Val Arg Lys Phe Asp Pro Met His Phe Lys Ser Phe Asp	225	230	235
10	His Leu Asn Ser Thr Ser Val Asn Glu Thr Glu Thr Thr Val Ala Thr	245	250	255
	Trp Gln Asp Gln Trp Glu Asp Val Ile Glu Arg Lys Ala Arg Ser Arg	260	265	270
	Arg Ala Ala Asn Ser Trp Asp His Tyr Val Glu Val Leu Val Val Ala	275	280	285
15	Asp Thr Lys Met Tyr Glu Tyr His Gly Arg Ser Leu Glu Asp Tyr Val	290	295	300
	Leu Thr Leu Phe Ser Thr Val Ala Ser Ile Tyr Arg His Gln Ser Leu	305	310	315
20	Arg Ala Ser Ile Asn Val Val Val Val Lys Leu Ile Val Leu Lys Thr	325	330	335
	Glu Asn Ala Gly Pro Arg Ile Thr Gln Asn Ala Gln Gln Thr Leu Gln	340	345	350
	Asp Phe Cys Arg Trp Gln Gln Tyr Tyr Asn Asp Pro Asp Asp Ser Ser	355	360	365
25	Val Gln His His Asp Val Ala Ile Leu Leu Thr Arg Lys Asp Ile Cys	370	375	380
	Arg Ser Gln Gly Lys Cys Asp Thr Leu Gly Leu Ala Glu Leu Gly Thr	385	390	395
30	Met Cys Asp Met Gln Lys Ser Cys Ala Ile Ile Glu Asp Asn Gly Leu	405	410	415

	Ser	Ala	Ala	Phe	Thr	Ile	Ala	His	Glu	Leu	Gly	His	Val	Phe	Ser	Ile	
				420					425					430			
	Pro	His	Asp	Asp	Glu	Arg	Lys	Cys	Ser	Thr	Tyr	Met	Pro	Val	Asn	Lys	
			435					440					445				
5	Asn	Asn	Phe	His	Ile	Met	Ala	Pro	Thr	Leu	Glu	Tyr	Asn	Thr	His	Pro	
			450				455					460					
	Trp	Ser	Trp	Ser	Pro	Cys	Ser	Ala	Gly	Met	Leu	Glu	Arg	Phe	Leu	Glu	
	465					470					475					480	
10	Asn	Asn	Arg	Gly	Gln	Thr	Gln	Cys	Leu	Phe	Asp	Gln	Pro	Val	Glu	Arg	
				485						490					495		
	Arg	Tyr	Tyr	Glu	Asp	Val	Phe	Val	Arg	Asp	Glu	Pro	Gly	Lys	Lys	Tyr	
				500					505					510			
	Asp	Ala	His	Gln	Gln	Cys	Lys	Phe	Val	Phe	Gly	Pro	Ala	Ser	Glu	Leu	
			515					520					525				
15	Cys	Pro	Tyr	Met	Pro	Thr	Cys	Arg	Arg	Leu	Trp	Cys	Ala	Thr	Phe	Tyr	
		530					535					540					
	Gly	Ser	Gln	Met	Gly	Cys	Arg	Thr	Gln	His	Met	Pro	Trp	Ala	Asp	Gly	
	545					550					555					560	
20	Thr	Pro	Cys	Asp	Glu	Ser	Arg	Ser	Met	Phe	Cys	His	His	Gly	Ala	Cys	
				565						570					575		
	Val	Arg	Leu	Ala	Pro	Glu	Ser	Leu	Thr	Lys	Ile	Asp	Gly	Gln	Trp	Gly	
				580					585					590			
	Asp	Trp	Arg	Ser	Trp	Gly	Glu	Cys	Ser	Arg	Thr	Cys	Gly	Gly	Gly	Val	
			595					600					605				
25	Gln	Lys	Gly	Leu	Arg	Asp	Cys	Asp	Ser	Pro	Lys	Pro	Arg	Asn	Gly	Gly	
		610					615					620					
	Lys	Tyr	Cys	Val	Gly	Gln	Arg	Glu	Arg	Tyr	Arg	Ser	Cys	Asn	Thr	Gln	
	625					630				635						640	
30	Glu	Cys	Pro	Trp	Asp	Thr	Gln	Pro	Tyr	Arg	Glu	Val	Gln	Cys	Ser	Glu	
				645						650					655		

	Phe	Asn	Asn	Lys	Asp	Ile	Gly	Ile	Gln	Gly	Val	Ala	Ser	Thr	Asn	Thr	
				660					665					670			
	His	Trp	Val	Pro	Lys	Tyr	Ala	Asn	Val	Ala	Pro	Asn	Glu	Arg	Cys	Lys	
			675					680					685				
5	Leu	Tyr	Cys	Arg	Leu	Ser	Gly	Ser	Ala	Ala	Phe	Tyr	Leu	Leu	Arg	Asp	
		690					695					700					
	Lys	Val	Val	Asp	Gly	Thr	Pro	Cys	Asp	Arg	Asn	Gly	Asp	Asp	Ile	Cys	
		705				710					715					720	
10	Val	Ala	Gly	Ala	Cys	Met	Pro	Ala	Gly	Cys	Asp	His	Gln	Leu	His	Ser	
					725					730					735		
	Thr	Leu	Arg	Arg	Asp	Lys	Cys	Gly	Val	Cys	Gly	Gly	Asp	Asp	Ser	Ser	
				740					745					750			
	Cys	Lys	Val	Val	Lys	Gly	Thr	Phe	Asn	Glu	Gln	Gly	Thr	Phe	Gly	Tyr	
			755					760					765				
15	Asn	Glu	Val	Met	Lys	Ile	Pro	Ala	Gly	Ser	Ala	Asn	Ile	Asp	Ile	Arg	
		770					775					780					
	Gln	Lys	Gly	Tyr	Asn	Asn	Met	Lys	Glu	Asp	Asp	Asn	Tyr	Leu	Ser	Leu	
		785				790					795					800	
20	Arg	Ala	Ala	Asn	Gly	Glu	Phe	Leu	Leu	Asn	Gly	His	Phe	Gln	Val	Ser	
				805						810					815		
	Leu	Ala	Arg	Gln	Gln	Ile	Ala	Phe	Gln	Asp	Thr	Val	Leu	Glu	Tyr	Ser	
				820					825					830			
	Gly	Ser	Asp	Ala	Ile	Ile	Glu	Arg	Ile	Asn	Gly	Thr	Gly	Pro	Ile	Arg	
			835					840					845				
25	Ser	Asp	Ile	Tyr	Val	His	Val	Leu	Ser	Val	Gly	Ser	His	Pro	Pro	Asp	
		850					855					860					
	Ile	Ser	Tyr	Glu	Tyr	Met	Thr	Ala	Ala	Val	Pro	Asn	Ala	Val	Ile	Arg	
		865				870					875					880	
30	Pro	Ile	Ser	Ser	Ala	Leu	Tyr	Leu	Trp	Arg	Val	Thr	Asp	Thr	Trp	Thr	
				885						890					895		

	Glu Cys Asp Arg Ala Cys Arg Gly Gln Gln Ser Gln Lys Leu Met Cys	
	900	905 910
	Leu Asp Met Ser Thr His Arg Gln Ser His Asp Arg Asn Cys Gln Asn	
	915	920 925
5	Val Leu Lys Pro Lys Gln Ala Thr Arg Met Cys Asn Ile Asp Cys Ser	
	930	935 940
	Thr Arg Trp Ile Thr Glu Asp Val Ser Ser Cys Ser Ala Lys Cys Gly	
	945	950 955 960
10	Ser Gly Gln Lys Arg Gln Arg Val Ser Cys Val Lys Met Glu Gly Asp	
	965	970 975
	Arg Gln Thr Pro Ala Ser Glu His Leu Cys Asp Arg Asn Ser Lys Pro	
	980	985 990
	Ser Asp Ile Ala Ser Cys Tyr Ile Asp Cys Ser Gly Arg Lys Trp Asn	
	995	1000 1005
15	Tyr Gly Glu Trp Thr Ser Cys Ser Glu Thr Cys Gly Ser Asn Gly Lys	
	1010	1015 1020
	Met His Arg Lys Ser Tyr Cys Val Asp Asp Ser Asn Arg Arg Val Asp	
	025	1030 1035 1040
20	Glu Ser Leu Cys Gly Arg Glu Gln Lys Glu Ala Thr Glu Arg Glu Cys	
	1045	1050 1055
	Asn Arg Ile Pro Cys Pro Arg Trp Val Tyr Gly His Trp Ser Glu Cys	
	1060	1065 1070
	Ser Arg Ser Cys Asp Gly Gly Val Lys Met Arg His Ala Gln Cys Leu	
	1075	1080 1085
25	Asp Ala Ala Asp Arg Glu Thr His Thr Ser Arg Cys Gly Pro Ala Gln	
	1090	1095 1100
	Thr Gln Glu His Cys Asn Glu His Ala Cys Thr Trp Trp Gln Phe Gly	
	1105	1110 1115 1120
30	Val Trp Ser Asp Cys Ser Ala Lys Cys Gly Asp Gly Val Gln Tyr Arg	
	1125	1130 1135

	Asp	Ala	Asn	Cys	Thr	Asp	Arg	His	Arg	Ser	Val	Leu	Pro	Glu	His	Arg	
				1140					1145					1150			
	Cys	Leu	Lys	Met	Glu	Lys	Ile	Ile	Thr	Lys	Pro	Cys	His	Arg	Glu	Ser	
		1155					1160						1165				
5	Cys	Pro	Lys	Tyr	Lys	Leu	Gly	Glu	Trp	Ser	Gln	Cys	Ser	Val	Ser	Cys	
		1170					1175					1180					
	Glu	Asp	Gly	Trp	Ser	Ser	Arg	Arg	Val	Ser	Cys	Val	Ser	Gly	Asn	Gly	
	185					1190					1195				1200		
10	Thr	Glu	Val	Asp	Met	Ser	Leu	Cys	Gly	Thr	Ala	Ser	Asp	Arg	Pro	Ala	
				1205						1210					1215		
	Ser	His	Gln	Thr	Cys	Asn	Leu	Gly	Thr	Cys	Pro	Phe	Trp	Arg	Asn	Thr	
				1220					1225					1230			
	Asp	Trp	Ser	Ala	Cys	Ser	Val	Ser	Cys	Gly	Ile	Gly	His	Arg	Glu	Arg	
		1235						1240					1245				
15	Thr	Thr	Glu	Cys	Ile	Tyr	Arg	Glu	Gln	Ser	Val	Asp	Ala	Ser	Phe	Cys	
		1250					1255					1260					
	Gly	Asp	Thr	Lys	Met	Pro	Glu	Thr	Ser	Gln	Thr	Cys	His	Leu	Leu	Pro	
	265				1270					1275					1280		
20	Cys	Thr	Ser	Trp	Lys	Pro	Ser	His	Trp	Ser	Pro	Cys	Ser	Val	Thr	Cys	
				1285						1290					1295		
	Gly	Ser	Gly	Ile	Gln	Thr	Arg	Ser	Val	Ser	Cys	Thr	Arg	Gly	Ser	Glu	
				1300					1305					1310			
	Gly	Thr	Ile	Val	Asp	Glu	Tyr	Phe	Cys	Asp	Arg	Asn	Thr	Arg	Pro	Arg	
			1315					1320					1325				
25	Leu	Lys	Lys	Thr	Cys	Glu	Lys	Asp	Thr	Cys	Asp	Gly	Pro	Arg	Val	Leu	
		1330					1335					1340					
	Gln	Lys	Leu	Gln	Ala	Asp	Val	Pro	Pro	Ile	Arg	Trp	Ala	Thr	Gly	Pro	
	1345				1350						1355				1360		
30	Trp	Thr	Ala	Cys	Ser	Ala	Thr	Cys	Gly	Asn	Gly	Thr	Gln	Arg	Arg	Leu	
				1365						1370					1375		

	Leu Lys Cys Arg Asp His Val Arg Asp Leu Pro Asp Glu Tyr Cys Asn	
	1380	1385 1390
	His Leu Asp Lys Glu Val Ser Thr Arg Asn Cys Arg Leu Arg Asp Cys	
	1395	1400 1405
5	Ser Tyr Trp Lys Met Ala Glu Trp Glu Glu Cys Pro Ala Thr Cys Gly	
	1410	1415 1420
	Thr His Val Gln Gln Ser Arg Asn Val Thr Cys Val Ser Ala Glu Asp	
	425	1430 1435 1440
10	Gly Gly Arg Thr Ile Leu Lys Asp Val Asp Cys Asp Val Gln Lys Arg	
	1445	1450 1455
	Pro Thr Ser Ala Arg Asn Cys Arg Leu Glu Pro Cys Pro Lys Gly Glu	
	1460	1465 1470
	Glu His Ile Gly Ser Trp Ile Ile Gly Asp Trp Ser Lys Cys Ser Ala	
	1475	1480 1485
15	Ser Cys Gly Gly Gly Trp Arg Arg Arg Ser Val Ser Cys Thr Ser Ser	
	1490	1495 1500
	Ser Cys Asp Glu Thr Arg Lys Pro Lys Met Phe Asp Lys Cys Asn Glu	
	505	1510 1515 1520
20	Glu Leu Cys Pro Pro Leu Thr Asn Asn Ser Trp Gln Ile Ser Pro Trp	
	1525	1530 1535
	Thr His Cys Ser Val Ser Cys Gly Gly Gly Val Gln Arg Arg Lys Ile	
	1540	1545 1550
	Trp Cys Glu Asp Val Leu Ser Gly Arg Lys Gln Asp Asp Ile Glu Cys	
	1555	1560 1565
25	Ser Glu Ile Lys Pro Arg Glu Gln Arg Asp Cys Glu Met Pro Pro Cys	
	1570	1575 1580
	Arg Ser His Tyr His Asn Lys Thr Ser Ser Ala Ser Met Thr Ser Leu	
	585	1590 1595 1600
30	Ser Ser Ser Asn Ser Asn Thr Thr Ser Ser Ala Ser Ala Ser Ser Leu	
	1605	1610 1615

	Pro	Ile	Leu	Pro	Pro	Val	Val	Ser	Trp	Gln	Thr	Ser	Ala	Trp	Ser	Ala	
				1620					1625					1630			
	Cys	Ser	Ala	Lys	Cys	Gly	Arg	Gly	Thr	Lys	Arg	Arg	Val	Val	Glu	Cys	
			1635					1640					1645				
5	Val	Asn	Pro	Ser	Leu	Asn	Val	Thr	Val	Ala	Ser	Thr	Glu	Cys	Asp	Gln	
		1650				1655						1660					
	Thr	Lys	Lys	Pro	Val	Glu	Glu	Val	Arg	Cys	Arg	Thr	Lys	His	Cys	Pro	
	665					1670					1675					1680	
10	Arg	Trp	Lys	Thr	Thr	Thr	Trp	Ser	Ser	Cys	Ser	Val	Thr	Cys	Gly	Arg	
				1685						1690					1695		
	Gly	Ile	Arg	Arg	Arg	Glu	Val	Gln	Cys	Tyr	Arg	Gly	Arg	Lys	Asn	Leu	
			1700						1705					1710			
	Val	Ser	Asp	Ser	Glu	Cys	Asn	Pro	Lys	Thr	Lys	Leu	Asn	Ser	Val	Ala	
		1715						1720					1725				
15	Asn	Cys	Phe	Pro	Val	Ala	Cys	Pro	Ala	Tyr	Arg	Trp	Asn	Val	Thr	Pro	
		1730					1735					1740					
	Trp	Ser	Lys	Cys	Lys	Asp	Glu	Cys	Ala	Arg	Gly	Gln	Lys	Gln	Thr	Arg	
	745					1750					1755					1760	
20	Arg	Val	His	Cys	Ile	Ser	Thr	Ser	Gly	Lys	Arg	Ala	Ala	Pro	Arg	Met	
				1765						1770					1775		
	Cys	Glu	Leu	Ala	Arg	Ala	Pro	Thr	Ser	Ile	Arg	Glu	Cys	Asp	Thr	Ser	
			1780					1785						1790			
	Asn	Cys	Pro	Tyr	Glu	Trp	Val	Pro	Gly	Asp	Trp	Gln	Thr	Cys	Ser	Lys	
		1795					1800						1805				
25	Ser	Cys	Gly	Glu	Gly	Val	Gln	Thr	Arg	Glu	Val	Arg	Cys	Arg	Arg	Lys	
		1810				1815						1820					
	Ile	Asn	Phe	Asn	Ser	Thr	Ile	Pro	Ile	Ile	Phe	Met	Leu	Glu	Asp	Glu	
	825					1830					1835					1840	
30	Pro	Ala	Val	Pro	Lys	Glu	Lys	Cys	Glu	Leu	Phe	Pro	Lys	Pro	Asn	Glu	
				1845						1850					1855		

	Ser	Gln	Thr	Cys	Glu	Leu	Asn	Pro	Cys	Asp	Ser	Glu	Phe	Lys	Trp	Ser	
				1860					1865					1870			
	Phe	Gly	Pro	Trp	Gly	Glu	Cys	Ser	Lys	Asn	Cys	Gly	Gln	Gly	Ile	Arg	
		1875						1880					1885				
5	Arg	Arg	Arg	Val	Lys	Cys	Val	Ala	Asn	Asp	Gly	Arg	Arg	Val	Glu	Arg	
		1890					1895					1900					
	Val	Lys	Cys	Thr	Thr	Lys	Lys	Pro	Arg	Arg	Thr	Gln	Tyr	Cys	Phe	Glu	
	905					1910					1915				1920		
10	Arg	Asn	Cys	Leu	Pro	Ser	Thr	Cys	Gln	Glu	Leu	Lys	Ser	Gln	Asn	Val	
					1925					1930					1935		
	Lys	Ala	Lys	Asp	Gly	Asn	Tyr	Thr	Ile	Leu	Leu	Asp	Gly	Phe	Thr	Ile	
			1940						1945					1950			
	Glu	Ile	Tyr	Cys	His	Arg	Met	Asn	Ser	Thr	Ile	Pro	Lys	Ala	Tyr	Leu	
		1955						1960					1965				
15	Asn	Val	Asn	Pro	Arg	Thr	Asn	Phe	Ala	Glu	Val	Tyr	Gly	Lys	Lys	Leu	
		1970					1975					1980					
	Ile	Tyr	Pro	His	Thr	Cys	Pro	Phe	Asn	Gly	Asp	Arg	Asn	Asp	Ser	Cys	
	985					1990					1995				2000		
20	His	Cys	Ser	Glu	Asp	Gly	Asp	Ala	Ser	Ala	Gly	Leu	Thr	Arg	Phe	Asn	
				2005						2010					2015		
	Lys	Val	Arg	Ile	Asp	Leu	Leu	Asn	Arg	Lys	Phe	His	Leu	Ala	Asp	Tyr	
				2020					2025					2030			
	Thr	Phe	Ala	Lys	Arg	Glu	Tyr	Gly	Val	His	Val	Pro	Tyr	Gly	Thr	Ala	
		2035					2040						2045				
25	Gly	Asp	Cys	Tyr	Ser	Met	Lys	Asp	Cys	Pro	Gln	Gly	Ile	Phe	Ser	Ile	
		2050					2055					2060					
	Asp	Leu	Lys	Ser	Ala	Gly	Leu	Lys	Leu	Val	Asp	Asp	Leu	Asn	Trp	Glu	
	065					2070				2075					2080		
30	Asp	Gln	Gly	His	Arg	Thr	Ser	Ser	Arg	Ile	Asp	Arg	Phe	Tyr	Asn	Asn	
				2085						2090					2095		

Ala Lys Val Ile Gly His Cys Gly Gly Phe Cys Gly Lys Cys Ser Pro
2100 2105 2110

Glu Arg Tyr Lys Gly Leu Ile Phe Glu Val Asn Thr Lys Leu Leu Asn
2115 2120 2125

5 His Val Lys Asn Gly Gly His Ile Asp Asp Glu Leu Asp Asp Asp Gly
2130 2135 2140

Phe Ser Gly Asp Met Asp
145 2150

10 QBMAD\195102